arXiv:1504.04083v1 [math.OA] 16 Apr 2015

THE COMPOSITION SERIES OF IDEALS OF THE PARTIAL-ISOMETRIC CROSSED PRODUCT BY SEMIGROUP OF ENDOMORPHISMS

SRIWULAN ADJI AND SAEID ZAHMATKESH

Abstract. Let Γ^+ be the positive cone in a totally ordered abelian group Γ, and α an action of Γ^+ by extendible endomorphisms of a C^*-algebra A. Suppose I is an extendible α-invariant ideal of A. We prove that the partial-isometric crossed product $\mathcal{I} := I \times_\alpha^{\text{piso}} \Gamma^+$ embeds naturally as an ideal of $A \times_\alpha^{\text{piso}} \Gamma^+$, such that the quotient is the partial-isometric crossed product of the quotient algebra. We claim that this ideal $\mathcal{I}$ together with the kernel of a natural homomorphism $\phi : A \times_\alpha^{\text{piso}} \Gamma^+ \to A \times_\alpha^{\text{iso}} \Gamma^+$ gives a composition series of ideals of $A \times_\alpha^{\text{piso}} \Gamma^+$ studied by Lindiarni and Raeburn.

1. Introduction

Let (A, Γ^+, α) be a dynamical system consisting of the positive cone Γ^+ in a totally ordered abelian group Γ, and an action $\alpha : \Gamma^+ \to \operatorname{End} A$ of Γ^+ by extendible endomorphisms of a C^*-algebra A. A covariant representation of the system (A, Γ^+, α) is defined for which the semigroup of endomorphisms $\{\alpha_s : s \in \Gamma^+\}$ are implemented by partial isometries, and then the associated partial-isometric crossed product C^*-algebra $A \times_\alpha^{\text{piso}} \Gamma^+$, generated by a universal covariant representation, is characterized by the property that its nondegenerate representations are in a bijective correspondence with covariant representations of the system. This generalizes the covariant isometric representation theory: the theory that uses isometries to represent the semigroup of endomorphisms in a covariant representation of the system. We denoted by $A \times_\alpha^{\text{iso}} \Gamma^+$ for the corresponding isometric crossed product.

Suppose I is an extendible α-invariant ideal of A, then $a + I \mapsto \alpha_x(a) + I$ defines an action of Γ^+ by extendible endomorphisms of the quotient algebra A/I. It is well-known that the isometric crossed product $I \times_\alpha^{\text{iso}} \Gamma^+$ sits naturally as an ideal in $A \times_\alpha^{\text{iso}} \Gamma^+$ such that $(A \times_\alpha^{\text{iso}} \Gamma^+)/(I \times_\alpha^{\text{iso}} \Gamma^+) \simeq A/I \times_\alpha^{\text{iso}} \Gamma^+$. We show that this result is valid for the partial-isometric crossed product.

Moreover if $\phi : A \times_\alpha^{\text{piso}} \Gamma^+ \to A \times_\alpha^{\text{iso}} \Gamma^+$ is the natural homomorphism given by the canonical universal covariant isometric representation of (A, Γ^+, α) in $A \times_\alpha^{\text{iso}} \Gamma^+$, then $\ker \phi$ together with the ideal $I \times_\alpha^{\text{piso}} \Gamma^+$ give a composition series of ideals of $A \times_\alpha^{\text{piso}} \Gamma^+$, from which we recover the structure theorems of [6]. Let us now consider the framework of [6]. A system that consists of the C^*-subalgebra $A := B_{\Gamma^+}$ of $\ell^\infty(\Gamma^+)$

2010 *Mathematics Subject Classification.* Primary 46L55.

Key words and phrases. C^*-algebra, endomorphism, semigroup, partial isometry, crossed product, primitive ideal, hull-kernel closure.

spanned by the functions 1_s satisfying

$$1_s(t) = \begin{cases} 1 & \text{if } t \geq s \\ 0 & \text{otherwise,} \end{cases}$$

and the action $\tau : \Gamma^+ \to \operatorname{End} B_{\Gamma^+}$ given by the translation on $\ell^\infty(\Gamma^+)$. We choose an extendible τ-invariant ideal I to be the subalgebra $B_{\Gamma^+,\infty}$ spanned by $\{1_x - 1_y : x < y \in \Gamma^+\}$. Then the composition series of ideals of $B_{\Gamma^+} \times_\tau^{\mathrm{piso}} \Gamma^+$, that is given by the two ideals $\ker \phi$ and $B_{\Gamma^+,\infty} \times_\tau^{\mathrm{piso}} \Gamma^+$, produces the large commutative diagram in [6, Theorem 5.6]. This result shows that the commutative diagram in [6, Theorem 5.6] exists for any totally ordered abelian subgroup (not only for subgroups of $\mathbb{R}$), and that we understand clearly where the diagram comes from.

Next, if we consider a specific semigroup Γ^+ such as the additive semigroup $\mathbb{N}$ in the group of integers $\mathbb{Z}$, then the large commutative diagram gives a clearer information about the ideals structure of $\mathbf{c} \times_\tau^{\mathrm{piso}} \mathbb{N}$. We can identify that the left-hand and top exact sequences in diagram [6, Theorem 5.6] are indeed equivalent to the extension of the algebra $\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}_0))$ of compact operators on the Hilbert module $\ell^2(\mathbb{N}, \mathbf{c}_0)$ by $\mathcal{K}(\ell^2(\mathbb{N}))$ provided by the algebra $\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}))$ of compact operators on $\ell^2(\mathbb{N}, \mathbf{c})$. Moreover it is known that $\operatorname{Prim} \mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c})) \simeq \operatorname{Prim}(\mathcal{K}(\ell^2(\mathbb{N})) \otimes \mathbf{c}) \simeq \operatorname{Prim} \mathbf{c}$ is homeomorphic to $\mathbb{N} \cup \infty$. Together with a knowledge about the primitive ideal space of the Toeplitz C^*-algebra generated by the unilateral shift, our theorem on the composition series of ideals of $\mathbf{c} \times_\tau^{\mathrm{piso}} \mathbb{N}$ provides a complete description of the topology on the primitive ideal space of $\mathbf{c} \times_\tau^{\mathrm{piso}} \mathbb{N}$.

We begin with a section containing background material about the partial-isometric crossed product by semigroups of extendible endomorphisms. In Section 3, we prove the existence of a short exact sequence of partial-isometric crossed products, which generalizes [2, Theorem 2.2] of the semigroup $\mathbb{N}$. Then we consider this and the other natural exact sequence described earlier in [4], to get the composition series of ideals in $A \times_\alpha^{\mathrm{piso}} \Gamma^+$.

We proceed to Section 4 by applying our results in Section 3 to the distinguished system $(B_{\Gamma^+}, \Gamma^+, \tau)$ and the extendible τ-invariant ideal $B_{\Gamma^+,\infty}$ of B_{Γ^+}. It can be seen from our Proposition 4.1 that the large commutative diagram of [6, Theorem 5.6] remains valid for any subgroup Γ of a totally ordered abelian group. Finally in the last section we describe the topology of primitive ideal space of $\mathbf{c} \times_\tau^{\mathrm{piso}} \mathbb{N}$ by using this large diagram.

2. Preliminaries

A bounded operator V on a Hilbert space H is called an isometry if $\|V(h)\| = \|h\|$ for all $h \in H$, which is equivalent to $V^*V = 1$. A bounded operator V on a Hilbert space H is called a *partial isometry* if it is isometry on $(\ker V)^\perp$. This is equivalent to $VV^*V = V$. If V is a partial isometry then so is the adjoint V^*, where as for an isometry V, the adjoint V^* may not be an isometry unless V is unitary. Associated to a partial isometry V, there are two orthogonal projections V^*V and VV^* on the initial space $(\ker V)^\perp$ and on the range VH respectively. In a C^*-algebra A, an element $v \in A$ is called an isometry if $v^*v = 1$ and a partial isometry if $vv^*v = v$.

An *isometric representation* of Γ^+ on a Hilbert space H is a map $S : \Gamma^+ \to B(H)$ which satisfies $S_x := S(x)$ is an isometry, and $S_{x+y} = S_x S_y$ for all $x, y \in \Gamma^+$. So an isometric representation of $\mathbb{N}$ is determined by a single isometry S_1. Similarly a *partial-isometric representation* of Γ^+ on a Hilbert space H is a map $V : \Gamma^+ \to B(H)$ which satisfies $V_x := V(x)$ is a partial isometry, and $V_{x+y} = V_x V_y$ for all $x, y \in \Gamma^+$. Note that the product VW of two partial isometries V and W is a partial isometry precisely when V^*V commutes with WW^* [6, Proposition 2.1]. Thus a partial isometry V is called a *power partial isometry* if V^n is a partial isometry for every $n \in \mathbb{N}$, so a partial-isometric representation of $\mathbb{N}$ is determined by a single power partial isometry V_1. If V is a partial-isometric representation of Γ^+, then every $V_x V_x^*$ commutes with $V_t V_t^*$, and so does $V_x^* V_x$ with $V_t^* V_t$.

Now we consider a dynamical system (A, Γ^+, α) consisting of a C^*-algebra A, an action α of Γ^+ by endomorphisms of A such that $\alpha_0 = \mathrm{id}$. Because we deal with non unital C^*-algebras and non unital endomorphisms, we require every endomorphism α_x to be extendible to a strictly continuous endomorphism $\overline{\alpha}_x$ on the multiplier algebra $M(A)$ of A. This happens precisely when there exists an approximate identity (a_λ) in A and a projection $p_{\alpha_x} \in M(A)$ such that $\alpha_x(a_\lambda)$ converges strictly to p_{α_x} in $M(A)$.

Definition 2.1. A *covariant isometric representation* of (A, Γ^+, α) on a Hilbert space H is a pair (π, S) of a nondegenerate representation $\pi : A \to B(H)$ and an isometric representation of $S : \Gamma^+ \to B(H)$ such that $\pi(\alpha_x(a)) = S_x \pi(a) S_x^*$ for all $a \in A$ and $x \in \Gamma^+$.

An *isometric crossed product* of (A, Γ^+, α) is a triple (B, j_A, j_{Γ^+}) consisting of a C^*-algebra B, a canonical covariant isometric representation (j_A, j_{Γ^+}) in $M(B)$ which satisfies the following

(i) for every covariant isometric representation (π, S) of (A, Γ^+, α) on a Hilbert space H, there exists a nondegenerate representation $\pi \times S : B \to B(H)$ such that $(\pi \times S) \circ j_A = \pi$ and $(\overline{\pi \times S}) \circ j_{\Gamma^+} = S$; and

(ii) B is generated by $j_A(A) \cup j_{\Gamma^+}(\Gamma^+)$, we actually have

$$B = \overline{\mathrm{span}}\{j_{\Gamma^+}(x)^* j_A(a) j_{\Gamma^+}(y) : x, y \in \Gamma^+, a \in A\}.$$

Note that a given system (A, Γ^+, α) could have a covariant isometric representation (π, S) only with $\pi = 0$. In this case the isometric crossed product yields no information about the system. If a system admits a non trivial covariant representation, then the isometric crossed product does exist, and it is unique up to isomorphism: if there is such a covariant isometric representation (t_A, t_{Γ^+}) of (A, Γ^+, α) in a C^*-algebra C, then there is an isomorphism of C onto B which takes (t_A, t_{Γ^+}) into (j_A, j_{Γ^+}). Thus we write the isometric crossed product B as $A \times_\alpha^{\mathrm{iso}} \Gamma^+$.

The partial-isometric crossed product of (A, Γ^+, α) is defined in a similar fashion involving partial-isometries instead of isometries.

Definition 2.2. A *covariant partial-isometric representation* of (A, Γ^+, α) on a Hilbert space H is a pair (π, S) of a nondegenerate representation $\pi : A \to B(H)$ and a partial-isometric representation $S : \Gamma^+ \to B(H)$ of Γ^+ such that $\pi(\alpha_x(a)) = S_x \pi(a) S_x^*$ for all $a \in A$ and $x \in \Gamma^+$. See in the Remark 2.3 that this equation implies $S_x^* S_x \pi(a) = \pi(a) S_x^* S_x$ for $a \in A$ and $x \in \Gamma^+$. Moreover, [6, Lemma 4.2] shows that every (π, S) extends to a partial-isometric covariant representation $(\overline{\pi}, S)$ of $(M(A), \Gamma^+, \overline{\alpha})$, and the partial-isometric covariance is equivalent to $\pi(\alpha_x(a)) S_x = S_x \pi(a)$ and $S_x S_x^* = \overline{\pi}(\overline{\alpha}_x(1))$ for $a \in A$ and $x \in \Gamma^+$.

A *partial-isometric crossed product* of (A, Γ^+, α) is a triple (B, j_A, j_{Γ^+}) consisting of a C^*-algebra B, a canonical covariant partial-isometric representation (j_A, j_{Γ^+}) in $M(B)$ which satisfies the following

(i) for every covariant partial-isometric representation (π, S) of (A, Γ^+, α) on a Hilbert space H, there exists a nondegenerate representation $\pi \times S : B \to B(H)$ such that $(\pi \times S) \circ j_A = \pi$ and $(\overline{\pi \times S}) \circ j_{\Gamma^+} = S$; and

(ii) B is generated by $j_A(A) \cup j_{\Gamma^+}(\Gamma^+)$, we actually have

$$B = \overline{\operatorname{span}}\{j_{\Gamma^+}(x)^* j_A(a) j_{\Gamma^+}(y) : x, y \in \Gamma^+, a \in A\}.$$

Unlike the theory of isometric crossed product: every system (A, Γ^+, α) admits a non trivial covariant partial-isometric representation (π, S) with π faithful [6, Example 4.6]. In fact [6, Proposition 4.7] shows that a canonical covariant partial-isometric representation (j_A, j_{Γ^+}) of (A, Γ^+, α) exists in the Toeplitz algebra $\mathcal{T}_X$ associated to a discrete product system X of Hilbert bimodules over Γ^+, which (i) and (ii) are fulfilled, and it is universal: if there is such a covariant partial-isometric representation (t_A, t_{Γ^+}) of (A, Γ^+, α) in a C^*-algebra C that satisfies (i) and (ii), then there is an isomorphism of C onto B which takes (t_A, t_{Γ^+}) into (j_A, j_{Γ^+}). Thus we write the partial-isometric crossed product B as $A \times_\alpha^{\mathrm{piso}} \Gamma^+$.

Remark 2.3. Our special thanks go to B. Kwasniewski for showing us the proof arguments in this remark. Assuming (π, S) is covariant, then by C^*-norm equation we have $\|\pi(a)S_x^* - S_x^*\pi(\alpha_x(a))\| = 0$, therefore $\pi(a)S_x^* = S_x^*\pi(\alpha_x(a))$ for all $a \in A$ and $x \in \Gamma^+$, which means that $S_x\pi(a) = \pi(\alpha_x(a))S_x$ for all $a \in A$ and $x \in \Gamma^+$. So $S_x^*S_x\pi(a) = S_x^*\pi(\alpha_x(a))S_x = (\pi(\alpha_x(a^*))S_x)^*S_x = (S_x\pi(a^*))^*S_x = \pi(a)S_x^*S_x$.

3. THE SHORT EXACT SEQUENCE OF PARTIAL-ISOMETRIC CROSSED PRODUCTS

Theorem 3.1. *Suppose that* $(A \times_\alpha^{\mathrm{piso}} \Gamma^+, i_A, V)$ *is the partial-isometric crossed product of a dynamical system* (A, Γ^+, α)*, and* I *is an extendible* α*-invariant ideal of* A*. Then there is a short exact sequence*

$$\begin{tikzcd} 0 \arrow[r] & I \times_\alpha^{\mathrm{piso}} \Gamma^+ \arrow[r, "\mu"] & A \times_\alpha^{\mathrm{piso}} \Gamma^+ \arrow[r, "\gamma"] & A/I \times_{\tilde{\alpha}}^{\mathrm{piso}} \Gamma^+ \arrow[r] & 0, \end{tikzcd} \tag{3.1}$$

where μ *is an isomorphism of* $I \times_\alpha^{\mathrm{piso}} \Gamma^+$ *onto the ideal*

$$\mathcal{D} := \overline{\operatorname{span}}\{V_x^* i_A(i) V_y : i \in I, x, y \in \Gamma^+\} \text{ of } A \times_\alpha^{\mathrm{piso}} \Gamma^+.$$

If $q : A \to A/I$ *is the quotient map,* i_I, W *denote the maps* $I \to I \times_\alpha^{\mathrm{piso}} \Gamma^+$*,* $W : \Gamma^+ \to M(I \times_\alpha^{\mathrm{piso}} \Gamma^+)$*, and similarly for* $i_{A/I}, U$ *the maps* $A/I \to A/I \times_{\tilde{\alpha}}^{\mathrm{piso}} \Gamma^+$*,* $\Gamma^+ \to M(A/I \times_{\tilde{\alpha}}^{\mathrm{piso}} \Gamma^+)$*, then*

$$\mu \circ i_I = i_A|_I, \quad \overline{\mu} \circ W = V \quad \text{and} \quad \gamma \circ i_A = i_{A/I} \circ q, \quad \overline{\gamma} \circ V = U.$$

Proof. We make some minor adjustment to the proof of [1, Theorem 3.1] for partial isometries. First, to check that $\mathcal{D}$ is indeed an ideal of $A \times_\alpha^{\mathrm{piso}} \Gamma^+$. Let $\xi = V_x^* i_A(i) V_y \in \mathcal{D}$. Then $V_s^*\xi$ is trivially contained in $\mathcal{D}$, and computations below show that $i_A(a)\xi$ and $V_s\xi$ are all in $\mathcal{D}$ for $a \in A$ and $s \in \Gamma^+$:

$$\begin{aligned} i_A(a)\xi &= i_A(a)V_x^* i_A(i) V_y = (V_x i_A(a^*))^* i_A(i) V_y \\ &= (i_A(\alpha_x(a^*))V_x)^* i_A(i) V_y = V_x^* i_A(\alpha_x(a) i) V_y; \end{aligned}$$

$$\begin{aligned}V_s\xi =&V_sV_x^*i_A(i)V_y = V_s(V_s^*V_sV_x^*V_x)V_x^*i_A(i)V_y = V_sV_u^*V_uV_x^*i_A(i)V_y, \quad u:=\max\{s,x\}\\ =&(V_sV_s^*V_{u-s}^*)(V_{u-x}V_xV_x^*)i_A(i)V_y = V_{u-s}^*(V_uV_u^*V_uV_u^*)(V_{u-x}i_A(i))V_y\\ =&V_{u-s}^*V_uV_u^*i_A(\alpha_{u-x}(i))V_{u-x}V_y = V_{u-s}^*\overline{i}_A(\overline{\alpha}_u(1))i_A(\alpha_{u-x}(i))V_{u-x+y}.\end{aligned}$$

This ideal $\mathcal{D}$ gives us a nondegenerate homomorphism $\psi : A\times_\alpha^{\mathrm{piso}}\Gamma^+ \to M(D)$ which satisfies $\psi(\xi)d = \xi d$ for $\xi \in A\times_\alpha^{\mathrm{piso}}\Gamma^+$ and $d\in\mathcal{D}$. Let $j_I : I \xrightarrow{i_A} A\times_\alpha^{\mathrm{piso}}\Gamma^+ \xrightarrow{\psi} M(\mathcal{D})$, and $S : \Gamma^+ \xrightarrow{V} M(A\times_\alpha^{\mathrm{piso}}\Gamma^+) \xrightarrow{\overline{\psi}} M(\mathcal{D})$. We use extendibility of ideal I to show j_I is nondegenerate. Take an approximate identity (e_λ) for I, and let $\varphi : A\to M(I)$ be the homomorphism satisfying $\varphi(a)i = ai$ for $a\in A$ and $i\in I$. Then $i_A(\alpha_s(e_\lambda)i)$ converges in norm to $i_A(\overline{\varphi}(\overline{\alpha}_s(1_{M(A)}))i)$. However

$$i_A(\overline{\varphi}(\overline{\alpha}_s(1_{M(A)}))i) = \overline{i}_A(\overline{\alpha}_s(1_{M(A)}))i_A(i) = V_sV_s^*i_A(i).$$

So $i_A(\alpha_s(e_\lambda)i)$ converges in norm to $V_sV_s^*i_A(i)$. Since $j_I(e_\lambda)V_s^*i_A(i)V_t = V_s^*i_A(\alpha_s(e_\lambda)i)V_t$ by covariance, it follows that $j_I(e_\lambda)V_s^*i_A(i)V_t$ converges in norm to $V_s^*i_A(i)V_t$. We can similarly show that $V_s^*i_A(i)V_tj_I(e_\lambda)$ converges in norm to $V_s^*i_A(i)V_t$. Thus $j_I(e_\lambda)\to 1_{M(\mathcal{D})}$ strictly, and hence j_I is nondegenerate.

We claim that the triple $(\mathcal{D}, j_I, S)$ is a partial-isometric crossed product of (I,Γ^+,α). A routine computations show the covariance of (j_I, S) for (I,Γ^+,α). Suppose now (π, T) is a covariant representation of (I,Γ^+,α) on a Hilbert space H. Let $\rho : A \xrightarrow{\varphi} M(I) \xrightarrow{\overline{\pi}} B(H)$. Then by extendibility of ideal I, that is $\overline{\alpha|_I}\circ\varphi = \varphi\circ\alpha$, the pair (ρ, T) is a covariant representation of (A,Γ^+,α). The restriction $(\rho\times T)|_{\mathcal{D}}$ to $\mathcal{D}$ of $\rho\times T$ is a nondegenerate representation of $\mathcal{D}$ which satisfies the requirement $(\rho\times T)|_{\mathcal{D}}\circ j_I = \pi$ and $\overline{(\rho\times T)|_{\mathcal{D}}}\circ S = T$. Thus the triple $(\mathcal{D}, j_I, S)$ is a partial-isometric crossed product for (I,Γ^+,α), and we have the homomorphism $\mu = i_A|_I\times V$.

Next we show the exactness. Let Φ be a nondegenerate representation of $A\times_\alpha^{\mathrm{piso}}\Gamma^+$ with kernel $\mathcal{D}$. Since $I\subset \ker\Phi\circ i_A$, we can have a representation $\tilde{\Phi}$ of A/I, which together with $\overline{\Phi}\circ V$ is a covariant partial-isometric representation of $(A/I,\Gamma^+,\tilde{\alpha})$. Then $\tilde{\Phi}\times(\overline{\Phi}\circ V)$ lifts to Φ, and therefore $\ker\gamma\subset\ker\Phi = \mathcal{D}$. □

Corollary 3.2. *Let (A,Γ^+,α) be a dynamical system, and I an extendible α-invariant ideal of A. Then there is a commutative diagram*

$$\begin{tikzcd}
 & 0 \arrow[d] & 0 \arrow[d] & 0 \arrow[d] & \\
0 \arrow[r] & \ker\phi_I \arrow[r] \arrow[d] & I\times_\alpha^{\mathrm{piso}}\Gamma^+ \arrow[r, "\phi_I"] \arrow[d, "\mu"'] & I\times_\alpha^{\mathrm{iso}}\Gamma^+ \arrow[r] \arrow[d, "\mu^{\mathrm{iso}}"'] & 0\\
0 \arrow[r] & \ker\phi_A \arrow[r] \arrow[d] & A\times_\alpha^{\mathrm{piso}}\Gamma^+ \arrow[r, "\phi_A"] \arrow[d, "\gamma"'] & A\times_\alpha^{\mathrm{iso}}\Gamma^+ \arrow[r] \arrow[d, "\gamma^{\mathrm{iso}}"'] & 0\\
0 \arrow[r] & \ker\phi_{A/I} \arrow[r] \arrow[d] & A/I\times_{\tilde{\alpha}}^{\mathrm{piso}}\Gamma^+ \arrow[r, "\phi_{A/I}"] \arrow[d] & A/I\times_{\tilde{\alpha}}^{\mathrm{iso}}\Gamma^+ \arrow[r] \arrow[d] & 0\\
 & 0 & 0 & 0 &
\end{tikzcd}$$

Proof. The three row exact sequences follow from [4], the middle column from Theorem 3.1 and the right column exact sequence from [1]. By inspection on the spanning elements, one can see that $\mu(\ker\phi_I)$ is an ideal of $\ker\phi_A$ and $\mu^{\mathrm{iso}}\circ\phi_I=\phi_A\circ\mu$, thus first and second rows commute. Then Snake Lemma gives the commutativity of all rows and columns. $\square$

4. The Example

We consider a dynamical system $(B_{\Gamma^+},\Gamma^+,\tau)$ consisting of a unital C^*-subalgebra B_{Γ^+} of $\ell^\infty(\Gamma^+)$ spanned by the set $\{1_s : s\in\Gamma^+\}$ of characteristic functions 1_s of $\{x\in\Gamma^+ : x\geq s\}$, the action τ of Γ^+ on B_{Γ^+} is given by $\tau_x(1_s)=1_{s+x}$. The ideal $B_{\Gamma^+,\infty}=\overline{\operatorname{span}}\{1_i-1_j : i<j\in\Gamma^+\}$ is an extendible τ-invariant ideal of B_{Γ^+}. Then we want to show in Proposition 4.1 that an application of Corollary 3.2 to the system $(B_{\Gamma^+},\Gamma^+,\tau)$ and the ideal $B_{\Gamma^+,\infty}$ gives [6, Theorem 5.6].

The crossed product $B_{\Gamma^+}\times_\tau^{\mathrm{iso}}\Gamma^+$ is a universal C^*-algebra generated by the canonical isometric representation t of Γ^+: every isometric representation w of Γ^+ gives a covariant isometric representation (π_w,w) of $(B_{\Gamma^+},\Gamma^+,\tau)$. Suppose $\{\varepsilon_x : x\in\Gamma^+\}$ is the usual orthonormal basis in $\ell^2(\Gamma^+)$, and let $T_s(\varepsilon_x)=\varepsilon_{x+s}$ for every $s\in\Gamma^+$. Then $s\mapsto T_s$ is an isometric representation of Γ^+, and the Toeplitz algebra $\mathcal{T}(\Gamma)$ is the C^*-subalgebra of $B(\ell^2(\Gamma^+))$ generated by $\{T_s : s\in\Gamma^+\}$. So there exists a representation $\mathfrak{T}:=\pi_T\times T$ of $B_{\Gamma^+}\times_\tau^{\mathrm{iso}}\Gamma^+$ on $\ell^2(\Gamma^+)$ such that $\mathfrak{T}(t_x)=T_x$ and $\mathfrak{T}(1_x)=T_xT_x^*$ for all $x\in\Gamma^+$. This representation is faithful by [3, Theorem 2.4]. Thus $B_{\Gamma^+}\times_\tau^{\mathrm{iso}}\Gamma^+$ and the Toeplitz algebra $\mathcal{T}(\Gamma)=\pi_T\times T(B_{\Gamma^+}\times_\tau^{\mathrm{iso}}\Gamma^+)$ are isomorphic, and the isomorphism takes the ideal $B_{\Gamma^+,\infty}\times_\tau^{\mathrm{iso}}\Gamma^+$ of $B_{\Gamma^+}\times_\tau^{\mathrm{iso}}\Gamma^+$ onto the commutator ideal $\mathcal{C}_\Gamma=\overline{\operatorname{span}}\{T_x(1-TT^*)T_y^* : x,y\in\Gamma^+\}$ of $\mathcal{T}(\Gamma)$.

Similarly, the crossed product $B_{\Gamma^+}\times_\tau^{\mathrm{piso}}\Gamma^+$ has a partial-isometric version of universal property by [6, Proposition 5.1]: every partial-isometric representation v of Γ^+ gives a covariant partial-isometric representation (π_v,v) of $(B_{\Gamma^+},\Gamma^+,\tau)$ with $\pi_v(1_x)=v_xv_x^*$, and then $B_{\Gamma^+}\times_\tau^{\mathrm{piso}}\Gamma^+$ is the universal C^*-algebra generated by the canonical partial-isometric representation v of Γ^+. Now since $x\mapsto T_x$ and $x\mapsto T_x^*$ are partial-isometric representations of Γ^+ in the Toeplitz algebra $\mathcal{T}(\Gamma)$, there exist (by the universality) a homomorphism φ_T and φ_{T^*} of $B_{\Gamma^+}\times_\tau^{\mathrm{piso}}\Gamma^+$ onto $\mathcal{T}(\Gamma)$.

Next consider the algebra $C(\hat{\Gamma})$ generated by $\{\lambda_x : x\in\Gamma\}$ of the evaluation maps $\lambda_x(\xi)=\xi(x)$ on $\hat{\Gamma}$. Let ψ_T and ψ_{T^*} be the homomorphisms of $\mathcal{T}(\Gamma)$ onto $C(\hat{\Gamma})$ defined by $\psi_T(T_x)=\lambda_x$ and $\psi_{T^*}(T_x)=\lambda_{-x}$.

Proposition 4.1. [6, Theorem 5.6] *Let Γ^+ be the positive cone in a totally ordered abelian group Γ. Then the following commutative diagram exists:*

$$\begin{tikzcd}
 & 0 \arrow[d] & 0 \arrow[d] & 0 \arrow[d] & \\
0 \arrow[r] & \ker\varphi_T \cap \ker\varphi_{T^*} \arrow[r] \arrow[d] & \ker\varphi_{T^*} \arrow[r, "\varphi_T|"] \arrow[d] & \mathcal{C}_\Gamma \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & \ker\varphi_T \arrow[r] \arrow[d] & B_{\Gamma^+} \times_\tau^{\mathrm{piso}} \Gamma^+ \arrow[r, "\varphi_T"] \arrow[d, "\varphi_{T^*}"'] \arrow[dr, "\Psi"] & \mathcal{T}(\Gamma) \arrow[r] \arrow[d, "\psi_{T^*}"] & 0 \\
0 \arrow[r] & \mathcal{C}_\Gamma \arrow[r] \arrow[d] & \mathcal{T}(\Gamma) \arrow[r, "\psi_T"] \arrow[d] & C(\hat\Gamma) \arrow[r] \arrow[d] & 0 \\
 & 0 & 0 & 0 &
\end{tikzcd}$$

(4.1)

where Ψ *maps each generator* $v_x \in B_{\Gamma^+} \times_\tau^{\mathrm{piso}} \Gamma^+$ *to* $\delta_x^* \in C^*(\Gamma) \simeq C(\hat\Gamma)$.

Proof. Apply Corollary 3.2 to the system $(B_{\Gamma^+}, \Gamma^+, \tau)$ and the extendible ideal $B_{\Gamma^+,\infty}$. Let $Q^{\mathrm{piso}} := B_{\Gamma^+}/B_{\Gamma^+,\infty} \times_{\tilde\tau}^{\mathrm{piso}} \Gamma^+$ and $Q^{\mathrm{iso}} := B_{\Gamma^+}/B_{\Gamma^+,\infty} \times_{\tilde\tau}^{\mathrm{iso}} \Gamma^+$. Then we have:

$$\begin{tikzcd}
 & 0 \arrow[d] & 0 \arrow[d] & 0 \arrow[d] & \\
0 \arrow[r] & \ker\phi_{B_{\Gamma^+,\infty}} \arrow[r] \arrow[d] & B_{\Gamma^+,\infty} \times_\tau^{\mathrm{piso}} \Gamma^+ \arrow[r, "\phi_{B_{\Gamma^+,\infty}}"] \arrow[d, "\mu"'] & B_{\Gamma^+,\infty} \times_\tau^{\mathrm{iso}} \Gamma^+ \arrow[r] \arrow[d, "\mu_{B_{\Gamma^+,\infty}}"] & 0 \\
0 \arrow[r] & \ker\phi_{B_{\Gamma^+}} \arrow[r] \arrow[d] & B_{\Gamma^+} \times_\tau^{\mathrm{piso}} \Gamma^+ \arrow[r, "\phi_{B_{\Gamma^+}}"] \arrow[d, "\gamma"'] & B_{\Gamma^+} \times_\tau^{\mathrm{iso}} \Gamma^+ \arrow[r] \arrow[d, "\gamma_{B_{\Gamma^+,\infty}}"] & 0 \\
0 \arrow[r] & \ker\phi_Q \arrow[r] \arrow[d] & Q^{\mathrm{piso}} \arrow[r, "\phi_Q"] \arrow[d] & Q^{\mathrm{iso}} \arrow[r] \arrow[d] & 0 \\
 & 0 & 0 & 0 &
\end{tikzcd}$$

(4.2)

We claim that exact sequences in this diagram and (4.1) are equivalent. The middle exact sequences of (4.1) and (4.2) are trivially equivalent via the isomorphism $\mathfrak{T} : B_{\Gamma^+} \times_\tau^{\mathrm{iso}} \Gamma^+ \to \mathcal{T}(\Gamma)$. By viewing B_{Γ^+} as the algebra of functions that have limit, the map $f \in B_{\Gamma^+} \mapsto \lim_{x \in \Gamma^+} f(x)$ induces an isomorphism $B_{\Gamma^+}/B_{\Gamma^+,\infty} \to \mathbb{C}$, which intertwines the action $\tilde\tau$ and the trivial action id on $\mathbb{C}$. So $(B_{\Gamma^+}/B_{\Gamma^+,\infty}, \Gamma^+, \tilde\tau) \simeq (\mathbb{C}, \Gamma^+, \mathrm{id})$. Moreover, $\mathfrak{T}$ combines with the isomorphism $h : B_{\Gamma^+}/B_{\Gamma^+,\infty} \times_{\tilde\tau}^{\mathrm{iso}} \Gamma^+ \to \mathbb{C} \times_{\mathrm{id}}^{\mathrm{iso}} \Gamma^+ \to C^*(\Gamma) \simeq C(\hat\Gamma)$ to identify the right-hand exact sequence equivalently to $0 \to \mathcal{C}_\Gamma \to \mathcal{T}(\Gamma) \overset{\psi_{T^*}}{\to} C(\hat\Gamma) \to 0$.

For the bottom sequence, we consider the pair of

$$\iota_{\mathbb{C}} : z \in \mathbb{C} \mapsto z1_{\mathcal{T}(\Gamma)} \text{ and } \iota_{\Gamma^+} : x \in \Gamma^+ \mapsto T_x^* \in \mathcal{T}(\Gamma).$$

It is a partial-isometric covariant representation, such that $(\mathcal{T}(\Gamma), \iota_{\mathbb{C}}, \iota_{\Gamma^+})$ is a partial-isometric crossed product of $(\mathbb{C}, \Gamma^+, \mathrm{id})$. So we have an isomorphism

$$\Upsilon : Q^{\mathrm{piso}} \to \mathbb{C} \times_{\mathrm{id}}^{\mathrm{piso}} \Gamma^+ \overset{\iota}{\to} \mathcal{T}(\Gamma) \text{ in which } \Upsilon(i_{\Gamma^+}(x)) = T_x^* \text{ for all } x,$$

and moreover if (j_Q, u) denotes the canonical covariant partial-isometric representation of the system $(Q := B_{\Gamma^+}/B_{\Gamma^+,\infty}, \Gamma^+, \tilde{\tau})$ in Q^{piso}, then Υ satisfies the equations $\Upsilon(u_x) = T_x^*$ and $\Upsilon(j_Q(1_x + B_{\Gamma^+,\infty})) = \iota_{\mathbb{C}}(\lim_y 1_x(y)) = 1$ for all $x \in \Gamma^+$. To see $\Upsilon(\ker \phi_Q) = \mathcal{C}_\Gamma$, recall from [4, Proposition 2.3] that

$$\ker \phi_Q := \overline{\operatorname{span}}\{u_x^* j_Q(a)(1 - u_z^* u_z)u_y : a \in Q, x, y, z \in \Gamma^+\}.$$

Since $\Upsilon(u_x^* j_Q(a)(1 - u_z^* u_z)u_y)$ is a scalar multiplication of $T_x(1 - T_z T_z^*)T_y^*$, therefore $\Upsilon(\ker \phi_Q) = \mathcal{C}_\Gamma$. Consequently the two exact sequences are equivalent:

$$\begin{tikzcd}
0 \arrow[r] & \ker \phi_Q \arrow[r] \arrow[d, "\Upsilon"'] & Q^{\mathrm{piso}} \arrow[r, "\phi_Q"] \arrow[d, "\Upsilon"'] & Q^{\mathrm{iso}} \arrow[r] \arrow[d, "h"'] & 0 \\
0 \arrow[r] & \mathcal{C}_\Gamma \arrow[r] & \mathcal{T}(\Gamma) \arrow[r, "\psi_T"] & C(\hat{\Gamma}) \arrow[r] & 0.
\end{tikzcd}$$

For the second column exact sequence, we note that the isomorphism $\jmath : Q^{\mathrm{piso}} \simeq \mathbb{C} \times_{\mathrm{id}}^{\mathrm{piso}} \Gamma^+ \to \mathcal{T}(\Gamma)$ satisfies $\jmath \circ \gamma = \varphi_{T^*}$. This implies

$$B_{\Gamma^+,\infty} \times_\tau^{\mathrm{piso}} \Gamma^+ \simeq \ker(\jmath \circ \gamma) = \ker \varphi_{T^*},$$

and therefore the second column sequence of diagram (4.1) is equivalent to $0 \to \ker \varphi_{T^*} \to B_{\Gamma^+} \times_\tau^{\mathrm{piso}} \Gamma^+ \to \mathcal{T}(\Gamma) \to 0$.

Next we are working for the first row. The homomorphism $\phi_{B_{\Gamma^+}}$ in the following diagram

$$\begin{tikzcd}
B_{\Gamma^+} \times_\tau^{\mathrm{piso}} \Gamma^+ \arrow[r, "\phi_{B_{\Gamma^+}}"] \arrow[dr, "\varphi_T"] & B_{\Gamma^+} \times_\tau^{\mathrm{iso}} \Gamma^+ \arrow[d, "\mathfrak{T}"'] \\
 & \mathcal{T}(\Gamma),
\end{tikzcd}$$

restricts to the homomorphism $\phi_{B_{\Gamma^+,\infty}}$ of the ideal $B_{\Gamma^+,\infty} \times_\tau^{\mathrm{piso}} \Gamma^+ \simeq \ker \varphi_{T^*}$ onto $B_{\Gamma^+,\infty} \times_\tau^{\mathrm{iso}} \Gamma^+ \simeq \mathcal{C}_\Gamma$. So the homomorphism $\varphi_T| : \ker \varphi_{T^*} \to \mathcal{C}_\Gamma$ has kernel $I := \ker \varphi_{T^*} \cap \ker \varphi_T$, and therefore first row exact sequence of the two diagrams are indeed equivalent.

Finally we show that such Ψ exists. Consider $C(\hat{\Gamma}) \simeq C^*(\Gamma) \simeq \mathbb{C} \times_{\mathrm{id}} \Gamma$ is the C^*-algebra generated by the unitary representation $x \in \Gamma \mapsto \delta_x \in \mathbb{C} \times_{\mathrm{id}} \Gamma$. Then we have a homomorphism $\pi_{\delta^*} \times \delta^* : B_{\Gamma^+} \times_\tau^{\mathrm{piso}} \Gamma^+ \to \mathbb{C} \times_{\mathrm{id}} \Gamma$ which satisfies $\pi_{\delta^*} \times \delta^*(v_x) = \delta_x^*$ for all $x \in \Gamma^+$, and hence it is surjective. By looking at the spanning elements of $\ker \varphi_T$ and $\ker \varphi_{T^*}$ we can see that these two ideals are contained in $\ker(\pi_{\delta^*} \times \delta^*)$, therefore $\mathcal{J} := \ker \varphi_T + \ker \varphi_{T^*}$ must be also in $\ker(\pi_{\delta^*} \times \delta^*)$. For the other inclusion, let ρ be a unital representation of $B_{\Gamma^+} \times_\tau^{\mathrm{piso}} \Gamma^+$ on a Hilbert space H_ρ with $\ker \rho = \mathcal{J}$. Then for $s \in \Gamma^+$ we have $\rho((1 - v_s v_s^*) - (1 - v_s^* v_s)) = 0$ because $1 - v_s v_s^* \in \ker \varphi_{T^*}$ and $1 - v_s^* v_s \in \ker \varphi_T$ belong to $\mathcal{J}$. So $0 = \rho(v_s^* v_s - v_s v_s^*)$, which implies that $\rho(v_s^* v_s) = \rho(v_s v_s^*)$. On the other hand the equation $\rho((1 - v_s v_s^*) + (1 - v_s^* v_s)) = 0$ gives $\rho(v_s v_s^*) = I$. Therefore $\rho(v_s v_s^*) = \rho(v_s^* v_s) = I$, and this means $\rho(v_s)$ is unitary for every $s \in \Gamma^+$. Consequently a representation $\tilde{\rho} : \mathbb{C} \times_{\mathrm{id}} \Gamma \to B(H_\rho)$ exists, and it satisfies $\tilde{\rho}\circ(\pi_{\delta^*}\times\delta^*)=\rho$. Thus $\ker\pi_{\delta^*}\times\delta^*\subset\ker\rho=\mathcal{J}$, and the composition $\pi_{\delta^*}\times\delta^*$ with the Fourier transform $C^*(\Gamma)\simeq C(\hat{\Gamma})$ is the wanted homomorphism Ψ. □

5. The Primitive Ideals of $\mathbf{c}\times_\tau^{\mathrm{piso}}\mathbb{N}$

Suppose Γ^+ is now the additive semigroup $\mathbb{N}$. The algebra $B_\mathbb{N}$ is conveniently viewed as the C^*-algebra $\mathbf{c}$ of convergent sequences, the ideal $B_{\mathbb{N},\infty}$ with $\mathbf{c}_0$, and the action τ of $\mathbb{N}$ on $\mathbf{c}$ is generated by the unilateral shift: $\tau_1(x_0,x_1,x_2,\cdots)=(0,x_0,x_1,x_2,\cdots)$. The universal C^*-algebra $\mathbf{c}\times_\tau^{\mathrm{piso}}\mathbb{N}$ is generated by a power partial isometry $v:=i_\mathbb{N}(1)$. The Toeplitz algebra $\mathcal{T}(\mathbb{Z})$ is the C^*-subalgebra of $B(\ell^2(\mathbb{N}))$ generated by isometries $\{T_n:n\in\mathbb{N}\}$, where $T_n(e_i)=e_{n+i}$ on the set of usual orthonormal basis $\{e_i:i\in\mathbb{N}\cup\{0\}\}$ of $\ell^2(\mathbb{N})$, and the commutator ideal of $\mathcal{T}(\mathbb{Z})$ is $\mathcal{K}(\ell^2(\mathbb{N}))$. Kernels of φ_T and φ_{T^*} are identified in [6, Lemma 6.2] by

$$\ker\varphi_T=\overline{\operatorname{span}}\{g^m_{i,j}:i,j,m\in\mathbb{N}\};\quad \ker\varphi_{T^*}=\overline{\operatorname{span}}\{f^m_{i,j}:i,j,m\in\mathbb{N}\}$$

where

$$g^m_{i,j}=v_i^*v_mv_m^*(1-v^*v)v_j\quad\text{and}\quad f^m_{i,j}=v_iv_m^*v_m(1-vv^*)v_j^*.$$

Moreover $\mathcal{I}:=\ker\varphi_T\cap\ker\varphi_{T^*}$ is an essential ideal in $\mathbf{c}\times_\tau^{\mathrm{piso}}\mathbb{N}$ [6, Lemma 6.8], given by

$$\overline{\operatorname{span}}\{f^m_{i,j}-f^{m+1}_{i,j}=g^m_{m-i,m-j}-g^{m+1}_{m-i,m-j}:m\in\mathbb{N},0\le i,j\le m\}.$$

The main point of [6, §6] is to show that there exist isomorphisms of $\ker\varphi_T$ and $\ker\varphi_{T^*}$ onto the algebra

$$\mathcal{A}:=\{f:\mathbb{N}\to K(\ell^2(\mathbb{N})):f(n)\in P_nK(\ell^2(\mathbb{N}))P_n\text{ and }\varepsilon_\infty(f)=\lim_n f(n)\text{ exists}\},$$

where $P_n:=1-T_{n+1}T^*_{n+1}$ is the projection of $\ell^2(\mathbb{N})$ onto the subspace spanned by $\{e_i:i=0,1,2,\cdots,n\}$, and such that they restrict to isomorphisms of $\mathcal{I}$ onto the ideal

$$\mathcal{A}_0:=\{f\in\mathcal{A}:\lim_n f(n)=0\}\text{ of }\mathcal{A}.$$

We shall show in Proposition 5.1 that $\mathcal{A}$ and $\mathcal{A}_0$ are related to the algebras of compact operators on the Hilbert $\mathbf{c}$-module $\ell^2(\mathbb{N},\mathbf{c})$ and on the closed sub-$\mathbf{c}$-module $\ell^2(\mathbb{N},\mathbf{c}_0)$. We supply our readers with some basic theory of the C^*-algebra of operators on this Hilbert module, and let us begin with recalling the module structure of $\ell^2(\mathbb{N},\mathbf{c})$ (and its closed sub-module). The vector space $\ell^2(\mathbb{N},\mathbf{c})$, containing all $\mathbf{c}$-valued functions $\mathsf{a}:\mathbb{N}\to\mathbf{c}$ such that the series $\sum_{n\in\mathbb{N}}\mathsf{a}(n)^*\mathsf{a}(n)$ converges in the norm of $\mathbf{c}$, forms a Hilbert $\mathbf{c}$-module with the module structure defined by $(\mathsf{a}\cdot x)(n)=\mathsf{a}(n)x$ for $x\in\mathbf{c}$, and its $\mathbf{c}$-valued inner product given by $\langle\mathsf{a},\mathsf{b}\rangle=\sum_{n\in\mathbb{N}}\mathsf{a}(n)^*\mathsf{b}(n)$. In fact the module $\ell^2(\mathbb{N},\mathbf{c})$ is naturally isomorphic to the Hilbert module $\ell^2(\mathbb{N})\otimes\mathbf{c}$ that arises from the completion of algebraic (vector space) tensor product $\ell^2(\mathbb{N})\odot\mathbf{c}$ associated to the $\mathbf{c}$-valued inner product defined on simple tensor product by $\langle\xi\otimes x,\eta\otimes y\rangle=\langle\xi,\eta\rangle x^*y$ for $\xi,\eta\in\ell^2(\mathbb{N})$ and $x,y\in\mathbf{c}$. The isomorphism is implemented by the map ϕ that takes $(e_i\otimes x)\in\ell^2(\mathbb{N})\otimes\mathbf{c}$ to the element $\phi(e_i\otimes x)\in\ell^2(\mathbb{N},\mathbf{c})$ which is the function $[\phi(e_i\otimes x)](n)=\begin{cases}x & \text{if } i=n\\ 0 & \text{otherwise.}\end{cases}$ By exactly the same arguments, we see that the two Hilbert $\mathbf{c}_0$-modules $\ell^2(\mathbb{N},\mathbf{c}_0)$ and $\ell^2(\mathbb{N})\otimes\mathbf{c}_0$ are isomorphic. However since $\mathbf{c}_0$ is an ideal of $\mathbf{c}$, it follows that the $\mathbf{c}_0$-module $\ell^2(\mathbb{N},\mathbf{c}_0)$ is a closed sub-$\mathbf{c}$-module of $\ell^2(\mathbb{N},\mathbf{c})$, and respectively $\ell^2(\mathbb{N}) \otimes \mathbf{c}_0$ is a closed sub-$\mathbf{c}$-module of $\ell^2(\mathbb{N}) \otimes \mathbf{c}$. Moreover the $\mathbf{c}$-module isomorphism ϕ restricts to $\mathbf{c}_0$-module isomorphism $\ell^2(\mathbb{N}, \mathbf{c}_0) \simeq \ell^2(\mathbb{N}) \otimes \mathbf{c}_0$.

Next, we consider the C^*-algebra $\mathcal{L}(\ell^2(\mathbb{N}, \mathbf{c}))$ of adjointable operators on $\ell^2(\mathbb{N}, \mathbf{c})$, and the ideal $\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}))$ of $\mathcal{L}(\ell^2(\mathbb{N}, \mathbf{c}))$ spanned by the set $\{\theta_{\mathsf{a},\mathsf{b}} : \mathsf{a}, \mathsf{b} \in \ell^2(\mathbb{N}, \mathbf{c})\}$ of compact operators on the module $\ell^2(\mathbb{N}, \mathbf{c})$. The algebra $\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}_0))$ is defined by the same arguments, and note that $\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}_0))$ is an ideal of $\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}))$. The isomorphism of two modules $\ell^2(\mathbb{N}, \mathbf{c})$ and $\ell^2(\mathbb{N}) \otimes \mathbf{c}$, implies that $\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c})) \simeq \mathcal{K}(\ell^2(\mathbb{N}) \otimes \mathbf{c})$, which by the Hilbert module theorem, this is the C^*-algebraic tensor product $\mathcal{K}(\ell^2(\mathbb{N})) \otimes \mathbf{c}$ of $\mathcal{K}(\ell^2(\mathbb{N}))$ and $\mathbf{c}$. We shall often use the characteristics functions $\{1_n : n \in \mathbb{N}\}$ as generator elements of $\mathbf{c}$ and the spanning set $\{\theta_{e_i \otimes 1_n, e_j \otimes 1_n} : i, j, n \in \mathbb{N}\}$ of $\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}))$ in our computations.

There is another ingredient that we need to consider to state the Proposition. Suppose $S \in \mathcal{L}(\ell^2(\mathbb{N}, \mathbf{c}))$ is an operator defined by $S(\mathsf{a})(i) = \mathsf{a}(i-1)$ for $i \geq 1$ and zero otherwise. One can see that $S^*S = 1$, i.e. S is an isometry. Let $p \in \mathcal{L}(\ell^2(\mathbb{N}, \mathbf{c}))$ be the projection $(p(\mathsf{a}))(n) = 1_n \mathsf{a}(n)$ for $\mathsf{a} \in \ell^2(\mathbb{N}, \mathbf{c})$, and similarly $q \in \mathcal{L}(\ell^2(\mathbb{N}, \mathbf{c}_0))$ be the projection $(q(\mathsf{a}))(n) = 1_n \mathsf{a}(n)$ for $\mathsf{a} \in \ell^2(\mathbb{N}, \mathbf{c}_0)$. Then the following two partial isometric representations of $\mathbb{N}$ in $p\mathcal{L}(\ell^2(\mathbb{N}, \mathbf{c}))p$ defined by

$$w : n \in \mathbb{N} \mapsto pS_n^* p \quad \text{and} \quad t : n \in \mathbb{N} \mapsto pS_n p,$$

induce the representations $\pi_w \times w$ and $\pi_t \times t$ of $\mathbf{c} \times_\tau^{\mathrm{piso}} \mathbb{N}$ in $p\mathcal{L}(\ell^2(\mathbb{N}, \mathbf{c}))p$ which satisfy $\pi_w \times w(v_i) = pS_i^* p$ and $\pi_t \times t(v_i) = pS_i p$ for all $i \in \mathbb{N}$. These $\pi_w \times w$ and $\pi_t \times t$ are faithful representations [4, Example 4.3].

Proposition 5.1. *The representations* $\pi_w \times w$ *and* $\pi_t \times t$ *map* $\ker \varphi_T$ *and* $\ker \varphi_{T^*}$ *isomorphically onto the full corner* $p\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}))p$*. Moreover, they restrict to isomorphisms of the ideal* $\ker \varphi_T \cap \ker \varphi_{T^*}$ *onto the full corner* $q\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}_0))q$*.*

Remark 5.2. It follows from this Proposition that $\operatorname{Prim} \ker \varphi_T$ and $\operatorname{Prim} \ker \varphi_{T^*}$ are both homeomorphic to $\operatorname{Prim} \mathbf{c}$. In fact, since $\ker \varphi_{T^*} \simeq \mathbf{c}_0 \times_\tau^{\mathrm{piso}} \mathbb{N}$ by [2, Corollary 3.1], we can therefore deduce that $\mathbf{c}_0 \times_\tau^{\mathrm{piso}} \mathbb{N}$ is Morita equivalent to $\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}))$. This is a useful fact for our subsequential work on the partial-isometric crossed product of lattice semigroup $\mathbb{N} \times \mathbb{N}$.

Proof of Proposition 5.1. We only have to show that $\pi_t \times t(\ker \varphi_{T^*}) = p\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}))p$ and $\pi_t \times t(\ker \varphi_T \cap \ker \varphi_{T^*}) = q\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}_0))q$. The rest of arguments is done in [4, Example 4.3].

Note that the algebra $p\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}))p$ is spanned by $\{p\theta^{\mathbf{c}}_{e_i \otimes 1_n, e_j \otimes 1_n} p : i, j, n \in \mathbb{N}\}$. Since $\pi_t \times t(f^n_{i,j}) = p\theta^{\mathbf{c}}_{e_i \otimes 1_n, e_j \otimes 1_n} p$ for every $i, j, n \in \mathbb{N}$, therefore $\pi_t \times t(\ker \varphi_{T^*}) = p\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}))p$.

Similarly we consider that $\{q\theta^{\mathbf{c}_0}_{e_i \otimes 1_{\{n\}}, e_j \otimes 1_{\{n\}}} q : i, j \leq n \in \mathbb{N}\}$ spans $q\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}_0))q$. We use the equation $\theta^{\mathbf{c}}_{e_i \otimes 1_n, e_j \otimes 1_0} = \theta^{\mathbf{c}}_{e_i \otimes 1_n, e_j \otimes 1_n}$ for every $n \in \mathbb{N}$, in the computations below, to see that

$$\begin{aligned}
\pi_t \times t(f^n_{i,j} - f^{n+1}_{i,j}) &= p(\theta^{\mathbf{c}}_{e_i \otimes 1_n, e_j \otimes 1_0} - \theta^{\mathbf{c}}_{e_i \otimes 1_{n+1}, e_j \otimes 1_0})p \\
&= p(\theta^{\mathbf{c}}_{(e_i \otimes 1_n) - (e_i \otimes 1_{n+1}), (e_j \otimes 1_0)})p = p(\theta^{\mathbf{c}}_{e_i \otimes 1_{\{n\}}, e_j \otimes 1_0})p \\
&= p(\theta^{\mathbf{c}}_{e_i \otimes 1_{\{n\}}, e_j \otimes 1_{\{n\}}})p.
\end{aligned}$$

To convince that every $p(\theta^{\mathbf{c}}_{e_i\otimes 1_{\{n\}},e_j\otimes 1_{\{n\}}})p$ belongs to $q\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}_0))q$, we need the embedding $\iota^{\mathcal{K}}$ of $q\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}_0))q$ in $p\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}))p$ stated in Lemma 5.3. In fact, every element $p(\theta^{\mathbf{c}}_{e_i\otimes 1_{\{n\}},e_j\otimes 1_{\{n\}}})p$ spans $\iota^{\mathcal{K}}(q\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}_0))q)$, therefore $\pi_t\times t(\ker\varphi_{T^*}\cap\ker\varphi_T) = \iota^{\mathcal{K}}(q\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}_0))q)$. □

Lemma 5.3. *Let* $p\in\mathcal{L}(\ell^2(\mathbb{N},\mathbf{c}))$ *and* $q\in\mathcal{L}(\ell^2(\mathbb{N},\mathbf{c}_0))$ *are the projections defined by* $(p(\mathsf{a}))(n) = 1_n\mathsf{a}(n)$ *for* $\mathsf{a}\in\ell^2(\mathbb{N},\mathbf{c})$, *and* $(q(\mathsf{a}))(n) = 1_n\mathsf{a}(n)$ *for* $\mathsf{a}\in\ell^2(\mathbb{N},\mathbf{c}_0)$. *Then the full corner* $q\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}_0))q$ *embeds naturally via* $\iota^{\mathcal{K}}(q\theta^{\mathbf{c}_0}_{\mathsf{a},\mathsf{b}}q) = p\theta^{\mathbf{c}}_{\mathsf{a},\mathsf{b}}p$ *as an ideal in* $p\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}))p$, *and there exists a short exact sequence*

$$0 \longrightarrow q\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}_0))q \xrightarrow{\iota^{\mathcal{K}}} p\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}))p \xrightarrow{q^{\mathcal{K}}} \mathcal{K}(\ell^2(\mathbb{N})) \longrightarrow 0,$$

where $q^{\mathcal{K}}(p\theta^{\mathbf{c}}_{\mathsf{a},\mathsf{b}}p) = \theta_{x,y}$ *with* $x,y\in\ell^2(\mathbb{N})$ *are given by* $x_i = \lim_{n\to\infty}(1_i\mathsf{a}(i))(n)$ *and* $y_i = \lim_{n\to\infty}(1_i\mathsf{b}(i))(n)$. *In particular we have*

$$\begin{aligned} q^{\mathcal{K}}(p\theta^{\mathbf{c}}_{e_i\otimes 1_n,e_j\otimes 1_m}p) = q^{\mathcal{K}}(\theta^{\mathbf{c}}_{p(e_i\otimes 1_n),p(e_j\otimes 1_m)}) &= q^{\mathcal{K}}(\theta^{\mathbf{c}}_{e_i\otimes 1_{n\vee i},e_j\otimes 1_{m\vee j}}) \\ &= T_i(1-TT^*)T_j^* \in \mathcal{K}(\ell^2(\mathbb{N})). \end{aligned}$$

Proof. Apply [5, Lemma 2.6] for the module $X := \ell^2(\mathbb{N},\mathbf{c})$ and $I = \mathbf{c}_0$. In this case we have the submodule $XI = \ell^2(\mathbb{N},\mathbf{c}_0)$. Note that if $\mathsf{a}\in\ell^2(\mathbb{N},\mathbf{c})$, then every sequence $\mathsf{a}(i)\in\mathbf{c}$ is convergent in $\mathbb{C}$, and the map $\mathsf{q}:\mathsf{a}\mapsto(\mathsf{q}(\mathsf{a}))(i) = \lim_{n\to\infty}(\mathsf{a}(i))(n)$ gives $0\to\ell^2(\mathbb{N},\mathbf{c}_0)\to\ell^2(\mathbb{N},\mathbf{c})\xrightarrow{\mathsf{q}}\ell^2(\mathbb{N})\to 0$. Moreover [5, Lemma 2.6] proves that $\iota^{\mathcal{K}}(\theta^{XI}_{\mathsf{a},\mathsf{b}}) = \theta^X_{\mathsf{a},\mathsf{b}}$ and $q^{\mathcal{K}}(\theta^X_{\mathsf{a},\mathsf{b}}) = \theta^{X/XI}_{\mathsf{q}(\mathsf{a}),\mathsf{q}(\mathsf{b})}$ give the exactness of the sequence

$$0 \longrightarrow \mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}_0)) \xrightarrow{\iota^{\mathcal{K}}} \mathcal{K}(\ell^2(\mathbb{N},\mathbf{c})) \xrightarrow{q^{\mathcal{K}}} \mathcal{K}(\ell^2(\mathbb{N})) \longrightarrow 0.$$

Since $\iota^{\mathcal{K}}(q\theta^{XI}_{\mathsf{a},\mathsf{b}}q) = \theta^X_{q(\mathsf{a}),q(\mathsf{b})} = p\theta^X_{\mathsf{a},\mathsf{b}}p$ for every a and b in $\ell^2(\mathbb{N},\mathbf{c}_0)$, the corner $q\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}_0))q$ is embedded into $p\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}))p$ such that $q^{\mathcal{K}}$ is defined by $q^{\mathcal{K}}(p\theta^X_{\mathsf{a},\mathsf{b}}p) = q^{\mathcal{K}}(\theta^X_{p(\mathsf{a}),p(\mathsf{b})}) = \theta_{x,y}$ where $x_i = \lim_{n\to\infty}(1_i\mathsf{a}(i))(n)$ and $y_i = \lim_{n\to\infty}(1_i\mathsf{b}(i))(n)$. Thus we obtain the required exact sequence. □

Proposition 5.4. *There are isomorphisms* $\Theta : p\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}))p\to\ker\varphi_T$ *and* $\Theta_* : p\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}))p\to\ker\varphi_{T^*}$ *defined by* $\Theta(p\theta^{\mathbf{c}}_{e_i\otimes 1_n,e_j\otimes 1_n}p) = g^n_{i,j}$ *and* $\Theta_*(p\theta^{\mathbf{c}}_{e_i\otimes 1_n,e_j\otimes 1_n}p) = f^n_{i,j}$ *for all* $i,j,n\in\mathbb{N}$ *such that the following commutative diagram has all rows and columns exact:*

$$\begin{tikzcd}
 & 0 \arrow[d] & 0 \arrow[d] & 0 \arrow[d] & \\
0 \arrow[r] & q\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}_0))q \arrow[r, "\iota^{\mathcal{K}}"] \arrow[d, "\iota^{\mathcal{K}} \circ \alpha"'] & p\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}))p \arrow[r, "q^{\mathcal{K}}"] \arrow[d, "\Theta_*"] & \mathcal{K}(\ell^2(\mathbb{N})) \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & p\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}))p \arrow[r, "\Theta"] \arrow[d, "q^{\mathcal{K}}"'] & \mathbf{c} \times_{\tau}^{\mathrm{piso}} \mathbb{N} \arrow[r, "\varphi_T"] \arrow[d, "\varphi_{T^*}"'] \arrow[dr, "\Psi"] & \mathcal{T}(\mathbb{Z}) \arrow[r] \arrow[d, "\psi_{T^*}"] & 0 \\
0 \arrow[r] & \mathcal{K}(\ell^2(\mathbb{N})) \arrow[r] \arrow[d] & \mathcal{T}(\mathbb{Z}) \arrow[r, "\psi_T"] \arrow[d] & C(\mathbb{T}) \arrow[r] \arrow[d] & 0 \\
 & 0 & 0 & 0 &
\end{tikzcd}$$

(5.1)

Proof. We apply Proposition 4.1 to the system $(\mathbf{c}, \mathbb{N}, \tau)$. Let $\{v_i : i \in \mathbb{N}\}$ denote the generators of $\mathbf{c} \times_{\tau}^{\mathrm{piso}} \mathbb{N}$, and $\{\delta_i : i \in \mathbb{Z}\}$ the generator of $C^*(\mathbb{Z})$. Then the homomorphism $\Psi : \mathbf{c} \times_{\tau}^{\mathrm{piso}} \mathbb{N} \to C(\mathbb{T})$ given by Proposition 4.1 satisfies $\Psi(v_i) = \delta_i^* = (z \mapsto \overline{z}^i) \in C(\mathbb{T})$ for every $i \in \mathbb{N}$. Moreover $\Theta = (\pi_w \times w)^{-1}$ and $\Theta_* = (\pi_t \times t)^{-1}$, by Proposition 5.1, satisfy $\Theta(p\theta^{\mathbf{c}}_{e_i \otimes 1_n, e_j \otimes 1_n} p) = g^n_{i,j}$ and $\Theta_*(p\theta^{\mathbf{c}}_{e_i \otimes 1_n, e_j \otimes 1_n} p) = f^n_{i,j}$ for all $i, j, n \in \mathbb{N}$. So the first row sequence is exact, and which is equivalent to the one of (4.1) for $\Gamma^+ = \mathbb{N}$ because

$$\begin{tikzcd}
0 \arrow[r] & \mathcal{I} \arrow[r, "\mathrm{id}"] \arrow[d, "\pi_t \times t"'] & \ker \varphi_{T^*} \arrow[r, "\varphi_T|"] \arrow[d, "\pi_t \times t"'] & \mathcal{K}(\ell^2(\mathbb{N})) \arrow[r] \arrow[d, "\mathrm{id}"'] & 0 \\
0 \arrow[r] & q\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}_0))q \arrow[r, "\iota^{\mathcal{K}}"] & p\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}))p \arrow[r, "q^{\mathcal{K}}"] & \mathcal{K}(\ell^2(\mathbb{N})) \arrow[r] & 0.
\end{tikzcd}$$

For the first column we use the automorphism α of $q\mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}_0))q$ defined on its spanning element by $\alpha(q\theta^{\mathbf{c}_0}_{e_i \otimes 1_{\{n\}}, e_j \otimes 1_{\{n\}}} q) = q\theta^{\mathbf{c}_0}_{e_{n-i} \otimes 1_{\{n\}}, e_{n-j} \otimes 1_{\{n\}}} q$. Then by inspections on the spanning elements of the algebras involved, we can see that the diagram (5.1) commutes. □

Thus we know from the diagram that the set $\operatorname{Prim} \mathbf{c} \times_{\tau}^{\mathrm{piso}} \mathbb{N}$ is given by the sets $\operatorname{Prim} \mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}))$ and $\operatorname{Prim} \mathcal{T}(\mathbb{Z})$. Since

$$\operatorname{Prim} \mathcal{T}(\mathbb{Z}) = \operatorname{Prim} \mathcal{K}(\ell^2(\mathbb{N})) \cup \operatorname{Prim} C(\mathbb{T}) = \{0\} \cup \mathbb{T},$$

and $\operatorname{Prim} \mathcal{K}(\ell^2(\mathbb{N},\mathbf{c}))$ is homeomorphic to

$$\operatorname{Prim} \mathbf{c} = \operatorname{Prim} \mathbf{c}_0 \cup \operatorname{Prim} \mathbb{C} \simeq \mathbb{N} \cup \{\infty\},$$

therefore $\operatorname{Prim} \mathbf{c} \times_{\tau}^{\mathrm{piso}} \mathbb{N}$ consists of a copy of $\{I_n\}$ of $\mathbb{N}$ embedded as an open subset, a copy of $\{J_z\}$ of $\mathbb{T}$ embedded as a closed subset. We identify these ideals in Proposition 5.7 and Lemma 5.12.

Note for now that $\ker \varphi_T$ and $\ker \varphi_{T^*}$ are primitive ideals of $\mathbf{c} \times_{\tau}^{\mathrm{piso}} \mathbb{N}$: the Toeplitz representation T of $\mathcal{T}(\mathbb{Z})$ on $\ell^2(\mathbb{N})$ is irreducible by [7, Theorem 3.13], and φ_T and φ_{T^*} are surjective homomorphisms of $\mathbf{c} \times_{\tau}^{\mathrm{piso}} \mathbb{N}$ onto $\mathcal{T}(\mathbb{Z})$, so $T \circ \varphi_T$ and $T \circ \varphi_{T^*}$

are irreducible representations of $\mathbf{c} \times_{\tau}^{\mathrm{piso}} \mathbb{N}$ on $\ell^2(\mathbb{N})$. Moreover, irreducibility of the representation $\mathrm{id} \circ q^{\mathcal{K}} : p\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}))p \xrightarrow{q^{\mathcal{K}}} \mathcal{K}(\ell^2(\mathbb{N})) \xrightarrow{\mathrm{id}} B(\ell^2(\mathbb{N}))$ implies the kernel $\mathcal{I} = \ker \varphi_T \cap \ker \varphi_{T^*} \simeq q\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}_0))q$ of $\mathrm{id} \circ q^{\mathcal{K}}$ is a primitive ideal of $p\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}))p \simeq \ker \varphi_T$. Similarly, $\mathcal{I}$ is a primitive ideal of $\ker \varphi_{T^*} \simeq p\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}))p$. Although $\mathcal{I} \notin \operatorname{Prim} \mathbf{c} \times_{\tau}^{\mathrm{piso}} \mathbb{N}$, the ideal $\mathcal{I}$ is essential in $\mathbf{c} \times_{\tau}^{\mathrm{piso}} \mathbb{N}$ by [6, Lemma 6.8], so the space $\operatorname{Prim} \mathcal{I} \simeq \operatorname{Prim} \mathbf{c}_0$ is dense in $\operatorname{Prim} \mathbf{c} \times_{\tau}^{\mathrm{piso}} \mathbb{N}$.

Next consider that $\mathcal{K}(\ell^2(\mathbb{N})) = \overline{\operatorname{span}}\{e_{ij} := T_i(1 - TT^*)T_j^* : i, j \in \mathbb{N}\}$, and recall that there is a natural isomorphism Λ of $\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c})) \simeq \mathcal{K}(\ell^2(\mathbb{N})) \otimes \mathbf{c}$ onto the algebra

$$C(\mathbb{N} \cup \{\infty\}, \mathcal{K}(\ell^2(\mathbb{N}))) := \{f : \mathbb{N} \to \mathcal{K}(\ell^2(\mathbb{N})) : \lim_n f(n) \text{ exists in } \mathcal{K}(\ell^2(\mathbb{N}))\}$$

given by $\Lambda(e_{ij} \otimes 1_k)(n) = 1_k(n)e_{ij}$ for $i, j, k, n \in \mathbb{N}$. Then $\Lambda(p\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}))p) \subset \mathcal{A}$ because

$$\begin{aligned}
[\Lambda(p(e_{ij} \otimes 1_m)p)](n) &= [\Lambda(e_{ij} \otimes 1_{m \vee i \vee j})](n) \\
&= \begin{cases} e_{ij} & \text{if } n \geq m \vee i \vee j \\ 0 & \text{otherwise} \end{cases} \\
&= \pi_n(f_{i,j}^m) = \pi_n^*(g_{i,j}^m).
\end{aligned}$$

Since $\Lambda = \pi \circ \Theta_* = \pi^* \circ \Theta$, Λ maps the corners $p\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}))p$ and $q\mathcal{K}(\ell^2(\mathbb{N}, \mathbf{c}_0))q$ isomorphically onto the algebra $\mathcal{A}$ and $\mathcal{A}_0$ respectively. Construction of this isomorphism in [6, §6] involves the representations π_n and π_n^*, for each $n \in \mathbb{N}$, of $\mathbf{c} \times_{\tau}^{\mathrm{piso}} \mathbb{N}$ on $\ell^2(\mathbb{N})$ that are associated to the partial-isometric representations $k \mapsto P_n T_k P_n$ and $k \mapsto P_n T_k^* P_n$ respectively, where $P_n := 1 - T_{n+1}T_{n+1}^*$ is the projection onto $H_n := \operatorname{span}\{e_i : i = 0, 1, 2, \cdots, n\}$. For every $a \in \ker \varphi_{T^*}$, the sequence $\{\pi_n(a)\}_{n \in \mathbb{N}}$ is convergent in $\mathcal{K}(\ell^2(\mathbb{N}))$, and then the map $a \in \ker \varphi_{T^*} \mapsto \pi(a) := \{\pi_n(a)\}_{n \in \mathbb{N}} \in \mathcal{A}$ defines the isomorphism.

These observations suggest that an extension of π should give a representation of $\mathbf{c} \times_{\tau}^{\mathrm{piso}} \mathbb{N}$ in the algebra $C_{\mathrm{b}}(\mathbb{N}, B(\ell^2(\mathbb{N})))$, and then primitive ideals are the kernels of evaluation maps. But we can consider a smaller algebra which gives more information on the image of π. Note that the algebra $C(\mathbb{N} \cup \{\infty\}, B(\ell^2(\mathbb{N})))$ is too small to consider, because the sequence $(P_n T_k P_n)_{n \in \mathbb{N}}$ as we see, does not converge to T_k in the operator norm on $B(\ell^2(\mathbb{N}))$, but it converges strongly to T_k. Therefore we consider the set $C_{\mathrm{b}}(\mathbb{N} \cup \{\infty\}, B(\ell^2(\mathbb{N}))_{*-\mathrm{s}})$ of functions $\xi : \mathbb{N} \to B(\ell^2(\mathbb{N}))$ such that $\lim_n \xi_n$ exists in the *-strong topology on $B(\ell^2(\mathbb{N}))$, and which satisfies $\|\xi\|_\infty := \sup_n \|\xi_n\| < \infty$. By [8, Lemma 2.56], it is a C^*-algebra with the pointwise operation from $B(\ell^2(\mathbb{N}))$ and the norm $\|\cdot\|_\infty$. Then let

$$\begin{aligned}
\mathcal{B} := \{f : \mathbb{N} \to B(\ell^2(\mathbb{N})) \; : \; &\sup_{n \in \mathbb{N}} \|f(n)\|_{B(\ell^2(\mathbb{N}))} < \infty, \; f(n) \in P_n B(\ell^2(\mathbb{N}))P_n \text{ and} \\
&\lim_{n \to \infty} f(n) \text{ exists in the } ^*\text{-strong topology on } B(\ell^2(\mathbb{N}))\}.
\end{aligned}$$

Note that $\mathcal{B}$ is a subalgebra of $C_{\mathrm{b}}(\mathbb{N} \cup \{\infty\}, B(\ell^2(\mathbb{N}))_{*-\mathrm{s}})$ because $P_n B(\ell^2(\mathbb{N}))P_n \simeq B(H_n)$ is closed in $B(\ell^2(\mathbb{N}))$ for every $n \in \mathbb{N}$, and $\mathcal{B}$ has an identity $1_{\mathcal{B}} = (P_0, P_1, P_2, \cdots)$.

Proposition 5.5. *There are faithful representations π and π^* of $\mathbf{c} \times_{\tau}^{\mathrm{piso}} \mathbb{N}$ in the algebra $\mathcal{B}$, which defined on each generator $v_k \in \mathbf{c} \times_{\tau}^{\mathrm{piso}} \mathbb{N}$ by*

$$\pi(v_k)(n) := \pi_n(v_k) = P_n T_k P_n \text{ and } \pi^*(v_k)(n) := \pi_n^*(v_k) = P_n T_k^* P_n \text{ for } n \in \mathbb{N}.$$

These representations π *and* π^* *are the extension of isomorphisms* $\pi : \ker \varphi_{T^*} \to \mathcal{A}$ *and* $\pi^* : \ker \varphi_T \to \mathcal{A}$ *of* [6, Theorem 6.1].

Proof. The map π is induced by the partial-isometric representation $k \mapsto W_k$ where $W_k(n) = P_n T_k P_n$, and similarly for π^* by $k \mapsto S_k$ where $S_k(n) = P_n T_k^* P_n$ for $n \in \mathbb{N}$. These are unital representations: $\pi(1) = \pi(v_0) = (P_0, P_1, P_2, \cdots) = \pi^*(1)$.

By [6, Proposition 5.4], the representation π is faithful if and only if for any $r > 0$ and $i < j$ in $\mathbb{N}$, we have $\xi^r_{i,j} \in \mathcal{B}$ for which

$$\xi^r_{i,j} := (\pi(1) - \pi(v_r)^*\pi(v_r))(\pi(v_i)\pi(v_i)^* - \pi(v_j)\pi(v_j)^*) \text{ is a nonzero element of } \mathcal{B}.$$

Let $r > 0$ and $i < j \in \mathbb{N}$, then we consider the three cases $0 < r \leq i < j$, $i < r < j$ and $i < j \leq r$ separately. If $0 < r \leq i < j$, then

$$\begin{aligned}
\xi^r_{i,j}(i) &= (P_i - \pi_i(v_r)^*\pi_i(v_r))(\pi_i(v_i)\pi_i(v_i)^* - \pi_i(v_j)\pi_i(v_j)^*) \\
&= (P_i - P_i T_r^* P_i T_r P_i)(P_i T_i P_i T_i^* P_i - P_i T_j P_i T_j^* P_i) \\
&= (P_i - P_i T_r^* T_r P_{i-r} P_i)(P_i T_i T_i^* P_i - 0) \\
&= (P_i - P_{i-r})(P_i T_i T_i^* P_i)
\end{aligned}$$

and that $[\xi^r_{i,j}(i)](e_i) = (P_i - P_{i-r})(e_i) = e_i$. If $i < j \leq r$ then similar computations show that $[\xi^r_{i,j}(i)](e_i) = [P_i(P_i T_i T_i^* P_i)](e_i) = e_i$, and for $i < r < j$ we have $[\xi^r_{i,j}(r)](e_r) = (P_r - P_0)(e_r) = e_r$. Thus $\xi^r_{i,j} \neq 0$ in $\mathcal{B}$. The same outline of arguments is valid to show the representation π^* is also faithful. $\square$

So we have for every $n \in \mathbb{N}$ the representations $\pi_n = \varepsilon_n \circ \pi$ and $\pi^*_n = \varepsilon_n \circ \pi^*$ of $\mathbf{c} \times^{\text{piso}}_{\tau} \mathbb{N}$ on H_n, where ε_n are the evaluation map of $C_{\text{b}}(\mathbb{N} \cup \{\infty\}, B(\ell^2(\mathbb{N}))_{*-\text{s}})$. Hence they are irreducible, indeed every nonzero vector of the subspace H_n of $\ell^2(\mathbb{N})$ is cyclic for π^*_n: if $(h_0, h_1, \cdots, h_n) \in H_n$ with $h_j \neq 0$ for some j, then for every $i \in \{0, 1, 2, \cdots, n\}$, we have

$$\begin{aligned}
(\pi^*_n(g^n_{i,j}))(h_0, h_1, \cdots, h_n) &= [T_i(1 - TT^*)T_j^*](h_0, h_1, \cdots, h_n) \\
&= (0, \cdots, h_j, \cdots, 0), \text{ where } h_j \text{ is in the } i\text{-th slot,}
\end{aligned}$$

so $\pi^*_n(\frac{1}{h_j}\ g^n_{i,j})(h) = e_i$, and therefore $H_n = \operatorname{span}\{\pi^*_n(\xi)h : \xi \in \mathbf{c} \times^{\text{piso}}_{\tau} \mathbb{N}\}$. Same arguments work for π_n.

Note for every $n \in \mathbb{N}$ that $\pi_n(f^m_{i,j}) = e_{ij} = \pi_n(g^k_{n-i,n-j})$ for all $0 \leq i, j, m, k \leq n$, and similarly $\pi^*_n(g^m_{i,j}) = e_{ij} = \pi^*_n(f^k_{n-i,n-j})$ for all $0 \leq i, j, m, k \leq n$. Thus every $f^m_{i,j} - g^k_{n-i,n-j}$ is contained in $\ker \pi_n$, and similarly $(g^m_{i,j} - f^k_{n-i,n-j}) \in \ker \pi^*_n$. We shall see many more elements of $\ker \pi_n$ as well as $\ker \pi^*_n$ in Proposition 5.7.

But now we recall that for $n \in \mathbb{N}$ the partial-isometric representation $J^n : \mathbb{N} \to B(H_n)$ in [6, §3] defined by $J^n_t(e_r) = \begin{cases} e_{t+r} & \text{if } r + t \in \{0, 1, \cdots, n\} \\ 0 & \text{otherwise,} \end{cases}$ induces the representation $\pi^{\mathbb{N}}_{J^n} \times J^n$ of $(\mathbf{c} \times^{\text{piso}}_{\tau} \mathbb{N}, v)$ on H_n. In fact $\pi^{\mathbb{N}}_{J^n} \times J^n = \pi_n$, because for every $k \in \mathbb{N}$ we have $(\pi^{\mathbb{N}}_{J^n} \times J^n(v_k))(e_r) = J^n_k(e_r) = P_n T_k P_n(e_r)$ where $r \in \{0, 1, 2, \cdots, n\}$.

The ideal $\ker \oplus^n_{r=0} \pi^{\mathbb{N}}_{J^r} \times J^r$ appears in the structure of $\mathbf{c} \times^{\text{piso}}_{\tau} \mathbb{N}$ [6, Lemma 5.7]. To be more precise about it, we need some results in [6, §5] related to the system $(\mathbb{C}^{n+1}, \tau, \mathbb{N})$. The crossed product $\mathbb{C}^{n+1} \times^{\text{piso}}_{\tau} \mathbb{N}$ is the universal C^*-algebra generated by a canonical partial-isometric representation w of $\mathbb{N}$ such that $w_r = 0$ for $r \geq n+1$. Let $q_n : (\mathbf{c} \times^{\text{piso}}_{\tau} \mathbb{N}, v) \to (\mathbb{C}^{n+1} \times^{\text{piso}}_{\tau} \mathbb{N}, w)$ be the homomorphism induced by w :

$\mathbb{N} \to \mathbb{C}^{n+1} \times_{\tau}^{\text{piso}} \mathbb{N}$, and note that it is surjective. Then Lemma 5.7 of [6] shows that $\ker q_n = \ker(\oplus_{r=0}^{n} \pi_{J^r}^{\mathbb{N}} \times J^r) = \bigcap_{r=0}^{n} \ker(\pi_{J^r}^{\mathbb{N}} \times J^r)$. So by these arguments we obtain the following equation

$$\ker q_n = \bigcap_{r=0}^{n} \ker \pi_r \text{ for every } n \in \mathbb{N}. \tag{5.2}$$

Lemma 5.6. *For $n \in \mathbb{N}$, let L_n be the ideal of $(\mathbf{c} \times_{\tau}^{\text{piso}} \mathbb{N}, v)$ generated by $\{v_r : r \geq n+1\}$. Then $L_n = \ker q_n$, and it is isomorphic to*

$$\{\xi \in \pi(\mathbf{c} \times_{\tau}^{\text{piso}} \mathbb{N}) \subset C_b(\mathbb{N} \cup \{\infty\}, B(\ell^2(\mathbb{N}))_{*-s}) : \xi \equiv 0 \text{ on } \{0, 1, 2, \cdots, n\}\}. \tag{5.3}$$

Proof. We have $L_n \subset \ker q_n$ because $q_n(v_k) = 0$ for all $k \geq n+1$. To see $\ker q_n \subset L_n$, let ρ be a representation of $\mathbf{c} \times_{\tau}^{\text{piso}} \mathbb{N}$ on H_ρ where $\ker \rho = L_n$. Since $\rho(v_t) = 0$ for every $t \geq n+1$, by the universal property of $\mathbb{C}^{n+1} \times_{\tau}^{\text{piso}} \mathbb{N}$, there exists a representation $\tilde{\rho}$ of $\mathbb{C}^{n+1} \times_{\tau}^{\text{piso}} \mathbb{N}$ on H_ρ which satisfies $\tilde{\rho} \circ q_n = \rho$. Thus $\ker q_n \subset \ker \rho = L_n$.

Next we show that $\pi(L_n)$ and (5.3) are equal. Let $r \geq n+1$, and consider $\pi(v_r)$ is the sequence $(P_i T_r P_i)_{i \in \mathbb{N}}$. If $0 \leq i \leq n$ then $0 \leq i+1 \leq n+1 \leq r$ and

$$P_i T_r P_i = (1 - T_{i+1} T_{i+1}^*) T_r P_i = (1 - T_{i+1} T_{i+1}^*) T_{i+1} T_{r-(i+1)} P_i = 0.$$

So $\pi(L_n)$ is a subset of (5.3). For the other inclusion, suppose $f \in \pi(\mathbf{c} \times_{\tau}^{\text{piso}} \mathbb{N})$ in which $f(i) = 0$ for all $0 \leq i \leq n$. Since $f = \pi(\xi)$ for some $\xi \in \mathbf{c} \times_{\tau}^{\text{piso}} \mathbb{N}$, and $\pi(\xi)(i) = \pi_i(\xi) = f(i)$ for all $i \in \mathbb{N}$, we therefore have $\pi_i(\xi) = f(i) = 0$ for all $0 \leq i \leq n$. Thus $\xi \in \cap_{i=0}^{n} \ker \pi_i = \ker q_n$, and hence $f = \pi(\xi) \in \pi(L_n)$. □

Let $\pi_\infty := \lim_n \pi_n$ and $\pi_\infty^* := \lim_n \pi_n^*$ where the limits are taken with respect to the strong topology of $B(\ell^2(\mathbb{N}))$. Then π_∞ and π_∞^* are the irreducible representations $\varphi_T : v_k \mapsto T_k$ and $\varphi_{T^*} : v_k \mapsto T_k^*$ of $\mathbf{c} \times_{\tau}^{\text{piso}} \mathbb{N}$ on $H_\infty := \ell^2(\mathbb{N})$. Thus by [6, Lemma 6.2] we have

$$\begin{aligned}\ker \pi_\infty &= \ker \varphi_T = \overline{\text{span}}\{g_{i,j}^m := v_i^* v_m v_m^* (1 - v^* v) v_j : i, j, m \in \mathbb{N}\} \\ \ker \pi_\infty^* &= \ker \varphi_{T^*} = \overline{\text{span}}\{f_{i,j}^m := v_i v_m^* v_m (1 - v v^*) v_j^* : i, j, m \in \mathbb{N}\}.\end{aligned}$$

For $n \in \mathbb{N}$, let π_n and π_n^* be the irreducible representations of $\mathbf{c} \times_{\tau}^{\text{piso}} \mathbb{N}$ on the subspace H_n of $\ell^2(\mathbb{N})$, that are induced by the partial-isometric representations $k \mapsto P_n T_k P_n$ and $k \mapsto P_n T_k^* P_n$. Let L_n be the ideal of $(\mathbf{c} \times_{\tau}^{\text{piso}} \mathbb{N}, v)$ generated by $\{v_r : r \geq n+1\}$. Then π_n is the representation

$$\varepsilon_n \circ \pi : \mathbf{c} \times_{\tau}^{\text{piso}} \mathbb{N} \xrightarrow{\pi} \mathcal{B} \subset C_{\text{b}}(\mathbb{N} \cup \{\infty\}, B(\ell^2(\mathbb{N}))_{*-\text{s}}) \xrightarrow{\varepsilon_n} B(H_n),$$

and similarly $\pi_n^* = \varepsilon_n \circ \pi^*$. So $\ker \pi_n \simeq \ker \varepsilon_n \simeq \ker \pi_n^*$.

Proposition 5.7. *Let $n \in \mathbb{N}$, then*

(a) $\ker \pi_n = \ker \pi_n^* \simeq \ker \varepsilon_n = \{\xi \in \mathcal{B} : \xi(n) = 0\}$*;*
(b) $\ker \pi_\infty \simeq \ker \pi_\infty^* = \{\xi \in \mathcal{B} : {}^*\!- \text{strong} \lim_n \xi(n) = 0\}$.

Furthermore

(c) $\ker \pi_n^* = \overline{\operatorname{span}}\{g^m_{i,j} - f^k_{n-i,n-j} + \eta : 0 \le i,j,m,k \le n, \eta \in L_n\}$,
$\ker \pi_n = \overline{\operatorname{span}}\{f^m_{i,j} - g^k_{n-i,n-j} + \eta : 0 \le i,j,m,k \le n, \eta \in L_n\}$, and
$\ker \pi_n^* = \ker \pi_n$ for $n \in \mathbb{N}$, in particular we have $\ker \pi_0 = \ker \pi_0^* = L_0$;

(d) $\ker \pi_n|_{\ker \varphi_{T^*}} = \overline{\operatorname{span}}\{(f^m_{i,j} - f^k_{i,j}) + f^z_{x,y} : 0 \le i,j,m,k \le n,$ one of $x,y,z \ge n+1\}$,
$\ker \pi_n^*|_{\ker \varphi_T} = \overline{\operatorname{span}}\{(g^m_{i,j} - g^k_{i,j}) + g^z_{x,y} : 0 \le i,j,m,k \le n,$ one of $x,y,z \ge n+1\}$,
$\Theta_*^{-1}(\ker \pi_n|_{\ker \varphi_{T^*}}) = \Theta^{-1}(\ker \pi_n^*|_{\ker \varphi_T})$, and
$\ker \pi_n^*|_{\ker \varphi_T} \simeq \{\mathsf{a} \in \mathcal{A} : \mathsf{a}(n) = 0\} \simeq \ker \pi_n|_{\ker \varphi_{T^*}}$;

(e) $\ker \pi_n^*|_{\mathcal{I}} = \overline{\operatorname{span}}\{g^m_{i,j} - g^{m+1}_{i,j} : 0 \le i,j \le m \text{ in } \mathbb{N}, \text{ and } m \ne n\} = \ker \pi_n|_{\mathcal{I}}$
$= \overline{\operatorname{span}}\{f^m_{i,j} - f^{m+1}_{i,j} : 0 \le i,j \le m \text{ in } \mathbb{N}, \text{ and } m \ne n\}$ is isomorphic to the ideal
$\{\mathsf{a} \in \mathcal{A}_0 : \mathsf{a}(n) = 0\}$.

Remark 5.8. Note that the representations $\pi_n|_{\ker \varphi_{T^*}}$ and $\pi_n^*|_{\ker \varphi_T}$ are equivalent to the evaluation map $\varepsilon_n : f \in \mathcal{A} \mapsto f(n) \in B(H_n)$ of $\mathcal{A}$ on H_n, so we have $\ker \pi_n|_{\ker \varphi_{T^*}} \simeq \ker \pi_n^*|_{\ker \varphi_T}$ is isomorphic to $\{f \in \mathcal{A} : f(n) = 0\}$, and $\ker \pi_n|_{\mathcal{I}} = \ker \pi_n^*|_{\mathcal{I}} \simeq \{f \in \mathcal{A}_0 : f(n) = 0\}$; and $\ker \pi_\infty \simeq \ker \pi_\infty^* \simeq \mathcal{A}$.

Proof of Proposition 5.7. Fix $n \in \mathbb{N}$. We show for $\ker \pi_n$, and skip the proof for $\ker \pi_n^*$ because it contains the same arguments. We clarify firstly that the space

$$\mathcal{J} := \overline{\operatorname{span}}\{f^m_{i,j} - g^k_{n-i,n-j} + \eta : 0 \le i,j,m,k \le n, \eta \in L_n\}$$

is an ideal of $(\mathbf{c} \times_\tau \mathbb{N}, v)$ by showing $v\mathcal{J} \subset \mathcal{J}$ and $v^*\mathcal{J} \subset \mathcal{J}$. Let $i = n$, then

$$\begin{aligned} vv_kv_k^*(1-v^*v)v_{n-j} = v(v^*vv_kv_k^*)(1-v^*v)v_{n-j} &= vv_kv_k^*v^*v(1-v^*v)v_{n-j} \\ &= v_{k+1}v_{k+1}^*(v - vv^*v)v_{n-j} = 0, \end{aligned}$$

therefore $v(f^m_{n,j} - g^k_{0,n-j} + \eta) = vv_nv_m^*v_m(1-vv^*)v_j^* - vv_kv_k^*(1-v^*v)v_{n-j} + v\eta = f^m_{n+1,j} + v\eta$ belongs to $\mathcal{J}$ because $f^m_{n+1,j} \in L_n$. If $0 \le i \le n-1$, then $1 \le i+1 \le n$ and $n - i \ge 1$, and we have

$$\begin{aligned} vv_{n-i}^*v_kv_k^* &= vv^*v_{n-i-1}^*v_{n-i-1}v_{n-i-1}^*v_kv_k^* \\ &= v_{n-i-1}^*v_{n-i-1}vv^*v_{n-i-1}^*v_kv_k^* \\ &= v_{n-i-1}^*v_{n-i}v_{n-i}^*v_kv_k^* = v_{n-i-1}^*v_{\max\{n-i,k\}}v_{\max\{n-i,k\}}^*, \end{aligned}$$

so $v(f^m_{i,j} - g^k_{n-i,n-j} + \eta) = f^m_{i+1,j} - g^{\max\{n-i,k\}}_{n-(i+1),n-j} + v\eta \in \mathcal{J}$.

Now we check for $v^*\mathcal{J}$, and assume $i = 0$, then

$$v^*[f^m_{0,j} - g^k_{n,n-j} + \eta] = v^*[v_m^*v_m(1-vv^*)v_j^* - v_n^*v_kv_k^*(1-v^*v)v_{n-j} + \eta] = 0 - g^k_{n+1,n-j} + v^*\eta \in \mathcal{J}$$

because $g^k_{n+1,n-j} \in L_n$. It follows by similar computations for $1 \le i \le n$ that

$$v^*[f^m_{i,j} - g^k_{n-i,n-j} + \eta] = f^{\max\{i,m\}}_{i-1,j} - g^k_{n-(i-1),n-j} + v^*\eta \in \mathcal{J}.$$

Next we show that $\mathcal{J} = \ker \pi_n$, one inclusion $\mathcal{J} \subset \ker \pi_n$ is clear because $\pi_n(f^m_{i,j}) = \pi_n(g^k_{n-i,n-j}) = T_i(1 - TT^*)T_j^*$ and $L_n \subset \ker \pi_n$. For the other inclusion, let $\sigma : \mathbf{c} \times_\tau^{\text{piso}} \mathbb{N} \to B(H_\sigma)$ be a nondegenerate representation with $\ker \sigma = \mathcal{J}$. Note that $B(H_n) = \operatorname{span}\{e_{ij} := T_i(1 - TT^*)T_j^* : 0 \le i,j \le n\}$. Since $\{f^n_{i,j} : 0 \le i,j \le n\}$ is a matrix-units for $B(H_\sigma)$, there is a homomorphism ψ of $B(H_n)$ into $B(H_\sigma)$ which satisfies $e_{ij} \mapsto \sigma(f^n_{i,j})$. Therefore $\sigma = \psi \circ \pi_n$, and hence $\ker \pi_n \subset \ker \sigma = \mathcal{J}$.

Using the spanning elements of $\ker \pi_n$ and $\ker \pi^*_n$, and the equation $f^m_{i,j} - g^k_{n-i,n-j} = -(g^k_{n-i,n-j} - f^m_{n-(n-i),n-(n-j)})$, we see that they contain each other, therefore $\ker \pi_n = \ker \pi^*_n$ for every $n \in \mathbb{N}$. The ideal L_0 is $\ker \pi_0 = \ker \pi^*_0$ because $f^0_{0,0} - g^0_{0,0} = v^*v - vv^* \in L_0$.

For (d), let now $\mathcal{J}$ be $\overline{\operatorname{span}}\{(f^m_{i,j} - f^k_{i,j}) + f^z_{x,y} : 0 \leq i, j, m, k \leq n,\ \text{one of } x, y, z \geq n+1\}$. Then the same idea of calculations shows that $\mathcal{J}$ is an ideal of $\ker \varphi_{T^*}$, and it is contained in $\ker \pi_n|_{\ker \varphi_{T^*}}$, then for the other inclusion let σ be a nondegenerate representation of $\ker \varphi_{T^*}$ such that $\ker \sigma = \mathcal{J}$, get the homomorphism $\psi : B(H_n) \to B(H_\sigma)$ defined by $\psi(e_{ij}) = \sigma(f^n_{i,j})$, and hence the equation $\psi \circ \pi_n = \sigma$ implies that $\ker \pi_n|_{\ker \varphi_{T^*}} = \mathcal{J}$. By computations on the spanning elements we see that the equation $\Theta^{-1}_*(\ker \pi_n|_{\ker \varphi_{T^*}}) = \Theta^{-1}(\ker \pi^*_n|_{\ker \varphi_T})$ is hold. The same arguments work for the proof of (e), and we skip this. □

Remark 5.9. The map $n \in \mathbb{N} \cup \{\infty\} \mapsto I_n := \ker \pi^*_n \in \operatorname{Prim}(\mathbf{c} \times^{\text{piso}}_\tau \mathbb{N})$ parameterizes the open subset $\{P \in \operatorname{Prim}(\mathbf{c} \times^{\text{piso}}_\tau \mathbb{N}) : \ker \varphi_T \simeq \mathcal{A} \not\subset P\}$ of $\operatorname{Prim}(\mathbf{c} \times^{\text{piso}}_\tau \mathbb{N})$ homeomorphic to $\operatorname{Prim} \mathcal{A}$. Note that the ∞ corresponds to the ideal $\ker \pi^*_\infty = \ker \varphi_{T^*} \in \operatorname{Prim}(\mathbf{c} \times^{\text{piso}}_\tau \mathbb{N})$, and it corresponds to $\mathcal{I} = \ker \varphi_{T^*}|_{\ker \varphi_T} \in \operatorname{Prim} \mathcal{A}$.

Lemma 5.10. (i) *$\bigcap_{n=0}^m I_n = L_m$ for every $m \in \mathbb{N}$;*

(ii) *$\bigcap_{n \in \mathbb{N}} I_n = \{0\}$;*

(iii) *$\{0\} \subsetneq (\bigcap_{n>m} I_n) \subset \ker \pi^*_\infty \cap \ker \pi_\infty$ for every $m \in \mathbb{N}$.*

Proof. Part (i) follows from (5.2) and Lemma 5.6. For (ii), note that q_∞ is the identity map on $\mathbf{c} \times^{\text{piso}}_\tau \mathbb{N}$, and that $\oplus_{i \in \mathbb{N}} \pi_i = (\oplus_{i \in \mathbb{N}}(\pi^{\mathbb{N}}_{J^i} \times J^i)) \circ \text{ id}$. So $\bigcap_{n \in \mathbb{N}} I_n = \{0\}$ by faithfulness of $\oplus_{i \in \mathbb{N}}(\pi^{\mathbb{N}}_{J^i} \times J^i)$ [6, Corollary 5.5].

The inclusion $\bigcap_{n>m} I_n \subset \ker \pi^*_\infty$ for every $m \in \mathbb{N}$ follows from the next arguments:

$$\begin{aligned}\bigcap_{n>m} \ker(\pi^*_n|_{\ker \pi_\infty}) &\simeq \{f \in \mathcal{A} : f(n) = 0\ \forall\ n > m\} \\ &\subset \{f \in A : \lim_{n \to \infty} f(n) = 0\} \\ &= \mathcal{A}_0 \simeq \ker \pi^*_\infty|_{\ker \pi_\infty} \subset \ker \pi^*_\infty \in \operatorname{Prim} \mathbf{c} \times^{\text{piso}}_\tau \mathbb{N},\end{aligned}$$

so the two ideals $J := \bigcap_{n>m} I_n$ and $L := \ker \pi_\infty$ of $\mathbf{c} \times^{\text{piso}}_\tau \mathbb{N}$ satisfy $J \cap L \subset \ker \pi^*_\infty$, therefore either $J \subset \ker \pi^*_\infty$ or $L \subset \ker \pi^*_\infty$, but the latter is not possible. To show $J \subset \ker \pi_\infty$, since $\ker \pi_n = \ker \pi^*_n$ for each n, we act similarly using the fact that

$$\bigcap_{n>m} \ker(\pi_n|_{\ker \pi^*_\infty}) \simeq \{f \in \mathcal{A} : f(n) = 0\ \forall\ n > m\} \subset \ker \pi_\infty \in \operatorname{Prim} \mathbf{c} \times^{\text{piso}}_\tau \mathbb{N}.$$

Therefore, $J \subset \ker \pi^*_\infty \cap \ker \pi_\infty$. Moreover, since $g^0_{0,0} - g^1_{0,0} \neq 0$ which satisfies $\pi^*_n(g^0_{0,0} - g^1_{0,0}) = 0$ for all $n \geq 1$, it follows that $\{0\} \subsetneq (\bigcap_{n>m} I_n)$. □

Remark 5.11. Part (ii) of Lemma 5.10 confirms with the fact that $\mathcal{I}$ is an essential ideal of $\mathbf{c} \times^{\text{piso}}_\tau \mathbb{N}$ [6, Lemma 6.8].

Next consider for $z \in \mathbb{T}$, the character $\gamma_z \in \hat{\mathbb{Z}} \simeq \mathbb{T}$ defined by $\gamma_z : m \mapsto \overline{z}^m$. Note that the map $\gamma_z : k \in \mathbb{N} \mapsto \gamma_z(k)$ is a partial-isometric representation of $\mathbb{N}$ in $\mathbb{C} \simeq B(\mathbb{C})$. Consequently for each $z \in \mathbb{T}$, we have a representation $\pi_{\gamma_z} \times \gamma_z$ of $\mathbf{c} \times^{\text{piso}}_\tau \mathbb{N}$

on $\mathbb{T}$ such that $\pi_{\gamma_z} \times \gamma_z(v_k) = \gamma_z(k) = \overline{z}^k$ for $k \in \mathbb{N}$, and it is irreducible. Moreover we know that the homomorphism $\Psi : \mathbf{c} \times_\tau^{\mathrm{piso}} \mathbb{N} \to C(\mathbb{T})$ is the composition of the Fourier transform $\mathbb{C} \times_{\mathrm{id}} \mathbb{Z} \simeq C^*(\mathbb{Z}) \simeq C(\mathbb{T})$ with $\ell \times \delta^* : \mathbf{c} \times_\tau^{\mathrm{piso}} \mathbb{N} \to \mathbb{C} \times_{\mathrm{id}} \mathbb{Z}$, in which $\ell : (x_n) \in \mathbf{c} \mapsto \lim_n x_n \in \mathbb{C}$ and δ is the unitary representation of $\mathbb{Z}$ on $\mathbb{C} \times_{\mathrm{id}} \mathbb{Z}$.

Lemma 5.12. *For $z \in \mathbb{T}$, the character $\gamma_z : k \mapsto \overline{z}^k$ in $\hat{\mathbb{Z}} \simeq \mathbb{T}$ gives an irreducible representation $\pi_{\gamma_z} \times \gamma_z$ of $\mathbf{c} \times_\tau^{\mathrm{piso}} \mathbb{N}$ on $\mathbb{T}$ such that $\pi_{\gamma_z} \times \gamma_z = \varepsilon_z \circ (\ell \times \delta^*)$. Denote by J_z the primitive ideal $\ker \pi_{\gamma_z} \times \gamma_z$ of $\mathbf{c} \times_\tau^{\mathrm{piso}} \mathbb{N}$. Then $\ker \pi_\infty$ and $\ker \pi_\infty^*$ are contained in J_z for every $z \in \mathbb{T}$. Moreover every ideal I_n for $n \in \mathbb{N}$ is not contained in any J_z.*

Proof. By using the Fourier transform we can view $\mathbb{C} \times_{\mathrm{id}} \mathbb{Z} \simeq C^*(\mathbb{Z})$ as $C(\mathbb{T})$, and it follows that $v_k \in \mathbf{c} \times_\tau^{\mathrm{piso}} \mathbb{N}$ is mapped into the function $\iota_k : t \mapsto \overline{t}^k \in C(\mathbb{T})$.

We know that primitive ideals of $C(\mathbb{T})$ are given by the kernels of evaluation maps $\varepsilon_t(f) = f(t)$ for $t \in \mathbb{T}$, and the character γ_z is a partial-isometric representation of $\mathbb{N}$ on $\mathbb{T}$ for $z \in \mathbb{T}$. Then by inspection on the generators, we see that the representation $\pi_{\gamma_z} \times \gamma_z$ of $\mathbf{c} \times_\tau^{\mathrm{piso}} \mathbb{N}$ on $\mathbb{T}$ satisfies $\pi_{\gamma_z} \times \gamma_z = \varepsilon_z \circ (\ell \times \delta^*)$. So the primitive ideal $J_z := \ker \pi_{\gamma_z} \times \gamma_z$ of $\mathbf{c} \times_\tau^{\mathrm{piso}} \mathbb{N}$ is lifted from the quotient $(\mathbf{c} \times_\tau^{\mathrm{piso}} \mathbb{N})/J \simeq C(\mathbb{T})$.

Since $\pi_{\gamma_z} \times \gamma_z(f_{i,j}^m) = 0 = \pi_{\gamma_z} \times \gamma_z(g_{i,j}^m)$, $\ker \pi_\infty = \ker \varphi_T$ and $\ker \pi_\infty^* = \ker \varphi_{T^*}$ are contained in J_z for every $z \in \mathbb{T}$. Finally, since $\pi_{\gamma_z} \times \gamma_z(v_{n+1}) = \overline{z}^{n+1} \neq 0$ for $n \in \mathbb{N}$, $I_n \not\subset J_z$ for any $z \in \mathbb{T}$. $\square$

Theorem 5.13. *The maps $n \in \mathbb{N} \cup \{\infty\} \cup \{\infty^*\} \mapsto I_n$ and $z \in \mathbb{T} \mapsto J_z$ combine to give a bijection of the disjoint union $\mathbb{N} \cup \{\infty\} \cup \{\infty^*\} \cup \mathbb{T}$ onto $\operatorname{Prim}(\mathbf{c} \times_\tau^{\mathrm{piso}} \mathbb{N})$, where $I_{\infty^*} := \ker \varphi_T$. Then the hull-kernel closure of a nonempty subset F of*

$$\mathbb{N} \cup \{\infty\} \cup \{\infty^*\} \cup \mathbb{T}$$

is given by

(a) *the usual closure of F in $\mathbb{T}$ if $F \subset \mathbb{T}$;*
(b) *F if F is a finite subset of $\mathbb{N}$;*
(c) *$F \cup \mathbb{T}$ if $F \subset (\{\infty\} \cup \{\infty^*\})$;*
(d) *$F \cup (\{\infty\} \cup \{\infty^*\} \cup \mathbb{T})$ if $F \neq \mathbb{N}$ is an infinite subset of $\mathbb{N}$;*
(e) *$\mathbb{N} \cup \{\infty\} \cup \{\infty^*\} \cup \mathbb{T}$ if $\mathbb{N} \subseteq F$.*

Proof. The diagram 5.1 together with Proposition 5.7 give a bijection map of $\mathbb{N} \cup \{\infty\} \cup \{\infty^*\} \cup \mathbb{T}$ onto $\operatorname{Prim}(\mathbf{c} \times_\tau^{\mathrm{piso}} \mathbb{N})$.

Lemma 5.10 (ii) gives the closure of the subset F in (e), and Lemma 5.10 (iii) gives the closure of the subset F in (d). If $F \subset (\{\infty\} \cup \{\infty^*\})$, then $\overline{F} = F \cup \mathbb{T}$ because $\ker \pi_\infty^*, \ker \pi_\infty \subset J_z$ for every $z \in \mathbb{T}$ by Lemma 5.12.

To see that $\overline{F} = F$ for a finite subset $F = \{n_1, n_2, \cdots, n_j\}$ of $\mathbb{N}$, we note that if an ideal $P \in \operatorname{Prim}(\mathbf{c} \times_\tau \mathbb{N})$ satisfies $\bigcap_{i=1}^{j} I_{n_i} \subset P$, then

- $P \neq J_z$ for any $z \in \mathbb{T}$ because $v_{n_j+1} \in \bigcap_{i=1}^{j} I_{n_i}$ but $v_{n_j+1} \notin J_z$;
- $P \neq I_\infty, I_{\infty^*}$ because $v_{n_j+1} \in \bigcap_{i=1}^{j} I_{n_i}$ but $v_{n_j+1} \notin I_\infty, I_{\infty^*}$;
- $P \neq I_n$ for $n \notin F$ because $(g_{0,0}^n - g_{0,0}^{n+1}) \in \bigcap_{i=1}^{j} I_{n_i}$ but $(g_{0,0}^n - g_{0,0}^{n+1}) \notin I_n$ for $n \notin F$.

So it can only be $P = I_j$ for some $j \in F$. Finally the usual closure of F in $\mathbb{T}$ is followed by the fact that the map $z \mapsto J_z$ is a homeomorphism of $\mathbb{T}$ onto the closed set $\operatorname{Prim} C(\mathbb{T})$. $\square$

References

[1] S. Adji, *Invariant ideals of crossed products by semigroups of endomorphisms*, Proc. Conference in Functional Analysis and Global Analysis in Manila, October 1996 (Springer, Singapore 1996), 1–8.

[2] S. Adji and A. Hosseini, *The Partial-Isometric Crossed Products of* $\mathbf{c}_0$ *by the Forward and the Backward Shifts*, Bull. Malays. Math. Sci. Soc. (2) **33**(3) (2010), 487–498.

[3] S. Adji, M. Laca, M. Nilsen and I. Raeburn, *Crossed products by semigroups of endomorphisms and the Toeplitz algebras of ordered groups*, Proc. Amer. Math. Soc. **122** (1994), no. 4, 1133–1141.

[4] S. Adji, S. Zahmatkesh, *Partial-isometric crossed products by semigroups of endomorphisms as full corners*, J. Aust. Math. Soc. **96** (2014), 145–166.

[5] N. J. Fowler, P. Muhly and I. Raeburn, *Representations of Cuntz-Pimsner Algebras*, Indiana University Math. J. **52** (2003), no. 3 , 569–605.

[6] J. Lindiarni and I. Raeburn, *Partial-isometric crossed products by semigroups of endomorphisms*, J. Operator Theory **52** (2004), 61–87.

[7] G.J. Murphy, *Ordered groups and Toeplitz algebras*, J. Operator Theory **18** (1987), 303–326.

[8] I. Raeburn and D. P. Williams, *Morita Equivalence and Continuous-Trace* C^**-Algebras*, Mathematical Surveys and Monographs, 60 (American Mathematical Society, Providence, RI, 1998).

E-mail address: `sriwulan.adji@gmail.com`

E-mail address: `zahmatkesh.s@gmail.com`